

3.9.7 Factors Influencing Emotions

Most of the physiological changes that occur during intense emotion, result from activation of the sympathetic division of the autonomic nervous system as it prepares the body for emergency action.

The sympathetic system is responsible for the following changes.

1. Blood pressure and heart rate increase
2. Respiration becomes more rapid
3. The pupils of the eye dilate.
4. Electrical resistance of the skin decreases.
5. Blood sugar level increases to provide more energy
6. The blood begins to clot more quickly in the case of wounds
7. Mobility of the gastro intestinal tract decreases or stops entirely. Blood is diverted from the stomach and intestines and sent to the brain and skeletal muscles.
8. The hairs on the skin erect causing a "Goose pimples. In emotion the sympathetic system also causes epinephrine (adrenaline) and non-epinephrine (noradrenalin). Nerve impulses with sympathetic system, which reach adrenal glands located on the top of the kidneys, trigger the secretion of hormones. They then get into the blood and circulate around the body.

Emotional stability and the nursing;

Nursing is a stressful job where traumatic situations are common. The ability to accept suffering and death without letting it gets personal as crucial.

That's not to say that there aren't heartwarming moments in nursing. Helping a patient to recover, reuniting families, or bonding with fellow nurses are special benefits of the job.

STUDENT'S ACTIVITY

- Let the student describe the feelings of joy, stress, and sad.
- Students can act out the emotions with facial expressions by creating face mask.

3.9.8 Motivation

It is the other factor which influence the behavior.

Definition - Motivation is an essential condition of learning process- **Melton**.

Nursing-voc_Unit 03.indd 71 22-02-2019 4.44.17 PM