Note

- The medians of a triangle are concurrent and the point of concurrence, the centroid G, is one-third of the distance from the opposite side to the vertex along the median.
- The centroid of the triangle obtained by joining the mid-points of the sides of a triangle is the same as the centroid of the original triangle.
- If (a_1, b_1), (a_2, b_2) and (a_3, b_3) are the mid-points of the sides of a triangle ABC then its centroid G is given by

$$G\left(\frac{a_1+a_2+a_3}{3}, \frac{b_1+b_2+b_3}{3}\right)$$




Fig. 5.39

FZB1QQ

Example 5.20 Find the centroid of the triangle whose veritices are $A(6,-1)$, $B(8,3)$ and $C(10,-5)$.

Solution

The centroid $G(x, y)$ of a triangle whose vertices are (x_1, y_1), (x_2, y_2) and (x_3, y_3) is given by

$$G(x, y) = G\left(\frac{x_1+x_2+x_3}{3}, \frac{y_1+y_2+y_3}{3}\right)$$

We have $(x_1, y_1) = (6,-1)$; $(x_2, y_2) = (8,3)$;
$(x_3, y_3) = (10,-5)$

The centroid of the triangle

$$G(x, y) = G\left(\frac{6+8+10}{3}, \frac{-1+3-5}{3}\right)$$

$$= G\left(\frac{24}{3}, \frac{-3}{3}\right) = G(8,-1)$$




Fig. 5.40

Note

- The Euler line of a triangle is the line that passes through the orthocenter (H), centroid (G) and the circumcenter (S). G divides the line segment $\overline{HS}$ in the ratio 2:1 from the orthocenter. That is centroid divides orthocenter and circumcenter internally in the ratio 2:1 from the Orthocentre.
- In an equilateral triangle, orthocentre, incentre, centroid and circumcentre are all the same.




Fig. 5.41

5-Coordinat_1.indd 216 26-12-2019 11:49:12