Points to remember

- If x_1, x_2 are the x-coordinates of two points on the x-axis then the distance between them is $x_2 - x_1$, if $x_2 > x_1$.
- If y_1, y_2 are the y-coordinates of two points on the y-axis then the distance between them is $|y_1 - y_2|$.
- Distance between the two points (x_1, y_1) and (x_2, y_2) is $\sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$
- Distance between (x_1, y_1) and the origin $(0, 0)$ is $\sqrt{x_1^2 + y_1^2}$
- The mid-point M of the line segment joining the points $A(x_1, y_1)$ and $B(x_2, y_2)$ is $M\left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2}\right)$
- The point P which divides the line segment joining the two points $A(x_1, y_1)$ and $B(x_2, y_2)$ internally in the ratio $m:n$ is $P\left(\frac{mx_2 + nx_1}{m+n}, \frac{my_2 + ny_1}{m+n}\right)$
- The centroid G of the triangle whose vertices are $A(x_1, y_1)$, $B(x_2, y_2)$ and $C(x_3, y_3)$ is $G\left(\frac{x_1 + x_2 + x_3}{3}, \frac{y_1 + y_2 + y_3}{3}\right)$
- The centroid of the triangle obtained by joining the mid-points of the sides of a triangle is the same as the centroid of the original triangle.

ICT Corner-1

Expected Result is shown in this picture



Step – 1

Open the Browser and copy and paste the Link given below (or) by typing the URL given (or) Scan the QR Code.

Step - 2

GeoGebra work book called "IX Analytical Geometry" will open. There are several worksheets given. Select the one you want. For example, open" Distance Formula"

Step-3

Move the sliders x_1, x_2, y_1, y_2 to change the co-ordinates of A and B. Now you calculate the distance AB using theDistance formula in a piece of paper and check your answer

B566_9_MAT_EM_T3

Browse in the link

Co-Ordinate Geometry: https://ggbm.at/V9HfY4v8

5-Coordinat_1.indd 220 26-12-2019 11:49:22