Mouse marking	1	2	3	4	5	6
Tumor Volume(mm^3)	145	148	142	141	139	140

Find the mean.

5. If the mean of the following data is 20.2, then find the value of p

Marks	10	15	20	25	30
No. of students	6	8	p	10	6

6. In the class, weight of students is measured for the class records. Calculate mean weight of the class students using Direct method.

Weight in kg	15-25	25-35	35-45	45-55	55-65	65-75
No. of students	4	11	19	14	0	2

7. Calculate the mean of the following distribution using Assumed Mean Method:

Class Interval	0-10	10-20	20-30	30-40	40-50
Frequency	5	7	15	28	8

8. Find the Arithmetic Mean of the following data using Step Deviation Method:

Age	15-19	20-24	25-29	30-34	35-39	40-44
No. of persons	4	20	38	24	10	9

8.5 Median

The arithmetic mean is typical of the data because it 'balances' the numbers; it is the number in the 'middle', pulled up by large values and pulled down by smaller values.

Suppose four people of an office have incomes of ₹5000, ₹6000, ₹7000 and ₹8000. Their mean income can be calculated as $\frac{5000+6000+7000+8000}{4}$ which gives ₹6500. If a fifth person with an income of ₹ 29000 is added to this group, then the arithmetic mean of all the five would be $\frac{5000+6000+7000+8000+29000}{5} = \frac{55000}{5}$ = ₹11000. Can one say that the average income of ₹11000 truly represents the income status of the individuals in the office? Is it not, misleading? The problem here is that an extreme score affects the Mean and can move the mean away from what would generally be considered the central area.

In such situations, we need a different type of average to provide reasonable answers.

8-Stats.indd 278 26-12-2019 14:18:40