When you search for some good video about Averages on You Tube, you look to watch the one with maximum views. Here you use the idea of a mode.

8.6.1 Mode - Raw Data

For an individual data mode is the value of the variable which occurs most frequently.

Example 8.14 In a rice mill, seven labours are receiving the daily wages of ₹500, ₹600, ₹600, ₹800, ₹800, ₹800 and ₹1000, find the modal wage.

Solution

In the given data ₹800 occurs thrice.Hence the mode is ₹ 800.

Example 8.15 Find the mode for the set of values 17, 18, 20, 20, 21, 21, 22, 22.

Solution

In this example, three values 20, 21, 22 occur two times each. There are three modes for the given data!

Note

- A distribution having only one mode is called unimodal.
- A distribution having two modes is called bimodal.
- A distribution having Three modes is called trimodal.
- A distribution having more than three modes is called multimodal.

8.6.2 Mode for Ungrouped Frequency Distribution

In a ungrouped frequency distribution, the value of the item having maximum frequency is taken as the mode.

Example 8.16 A set of numbers consists of five 4's, four 5's, nine 6's,and six 9's. What is the mode.

Solution

Size of item	4	5	6	9
Frequency	5	4	9	6

6 has the maximum frequency 9. Therefore 6 is the mode.

8-Stats.indd 285 26-12-2019 14:18:45