17. When $\left(2\sqrt{5}-\sqrt{2}\right)^2$ is simplified, we get

(1) $4\sqrt{5}+2\sqrt{2}$ (2) $22-4\sqrt{10}$ (3) $8-4\sqrt{10}$ (4) $2\sqrt{10}-2$

18. $\left(0.000729\right)^{\frac{-3}{4}} \times \left(0.09\right)^{\frac{-3}{4}}$ = _____

(1) $\frac{10^3}{3^3}$ (2) $\frac{10^5}{3^5}$ (3) $\frac{10^2}{3^2}$ (4) $\frac{10^6}{3^6}$

19. If $\sqrt{9^x} = \sqrt[3]{9^2}$, then x = _____

(1) $\frac{2}{3}$ (2) $\frac{4}{3}$ (3) $\frac{1}{3}$ (4) $\frac{5}{3}$

20. The length and breadth of a rectangular plot are 5×10^5 and 4×10^4 metres respectively. Its area is _____.

(1) $9\times10^1\ m^2$ (2) $9\times10^9\ m^2$ (3) $2\times10^{10}\ m^2$ (4) $20\times10^{20}\ m^2$

Points to Remember

- When the decimal expansion of $\frac{p}{q}$, q$\neq$0 terminates that is, comes to an end, the decimal is called a terminating decimal.
- In the decimal expansion of $\frac{p}{q}$, q$\neq$0 when the remainder is not zero, we have a repeating (recurring) block of digits in the quotient. In this case, the decimal expansion is called non-terminating and recurring.
- If a rational number $\frac{p}{q}$, q$\neq$ 0 can be expressed in the form $\frac{p}{2^m\times5^n}$, where $p\in\mathbb{Z}$ and $m, n\in\mathbb{W}$, then the rational number will have a terminating decimals. Otherwise, the rational number will have a non-terminating repeating (recurring) decimal.
- A rational number can be expressed either a terminating or a non- terminating recurring decimal.
- An irrational number is a non-terminating and non-recurring decimal, i.e. it cannot be written in form $\frac{p}{q}$, where p and q are both integers and q$\neq$ 0.
- The union of all rational numbers and all irrational numbers is called the set of real numbers.
- Every real number is either a rational number or an irratonal number.
- If a real number is not rational number, then it must be an irrational number.
- If 'a' is a positive rational number, 'n' is a positive integer and if $\sqrt[n]{a}$ is an irrational number, then $\sqrt[n]{a}$ is called as a surd.
- If 'm', 'n' are positive integers and a, b are positive rational numbers, then

(i) $\left(\sqrt[n]{a}\right)^n = a = \sqrt[n]{a^n}$ (ii) $\sqrt[n]{a}\times\sqrt[n]{b} = \sqrt[n]{ab}$ (iii) $\sqrt[m]{\sqrt[n]{a}} = \sqrt[mn]{a} = \sqrt[n]{\sqrt[m]{a}}$ (iv) $\frac{\sqrt[n]{a}}{\sqrt[n]{b}} = \sqrt[n]{\frac{a}{b}}$

- The process of multiplying a surd by another surd to get a rational number is called Rationalisation.
- Expressing a number N in the form of $N = a\times10^n$ where, $1\leq a<10$ and 'n' is an integer is called as Scientific Notation.

2-Real Numbers.indd 76 26-12-2019 13:35:46