Example

=ASIN (0) returns 0. =ASIN (1) returns 1.5707963267949 (PI/2 radians).

=DEGREES (ASIN (0.5)) returns 30. The sine of 30 degrees is 0.5.

ATAN (Number/Cell Address)

This function returns the inverse trigonometric tangent of Number where the angle is in radians. The angle returned is in the range -PI/2 to +PI/2. To return the angle in degrees, use the **DEGREES** function.

Example

=ATAN (1) returns 0.785398163397448 (PI/4 radians).

=DEGREES (ATAN (1)) returns 45. The tangent of 45 degrees is 1.

CEILING (Number; Significance; Mode)

This function rounds a number up to the nearest multiple of Significance. **Number** is the number that is to be rounded up. **Significance** is the value that the number is to be rounded up to a multiple of. **Mode** is an optional value. If the Mode parameter is supplied and is not equal to zero and if Number and Significance are negative, rounding up is carried out based on the absolute value of Number. This parameter is omitted when exporting to Microsoft Excel since Excel does not support a third parameter for this function

Example:

=CEILING (15.5;2;2) returns 16, =CEILING(-11;-2) returns -10

=CEILING (-11;-2;0) returns -10, =CEILING(-11;-2;1) returns -12

COMBIN (Count1; Count2)

Returns the number of combinations for a given number of objects (without repetition). Count1 is the number of items in the set. Count2 is the number of items to choose from the set. **COMBIN** returns the number of ordered ways to choose these items. For example if there are 3 items A, B and C in a set, you can choose 2 items in 3 different ways, namely AB, AC and BC. **COMBIN** implements the formula: Count1!/(Count2!*(Count1-Count2)!)

Example:

=COMBIN (3;2) returns 3, =COMBIN(5;3) returns 10.

COMBINA (Count1; Count2)

Returns the number of combinations of a subset of items including repetitions. **Count1** is the number of items in the set. **Count2** is the number of items to choose from the set. **COMBINA** returns the number of unique ways to choose these items, where the order of choosing is irrelevant, and repetition of items is allowed. For example if there are 3 items A, B and C in a set, you can choose 2 items in 6 different ways, namely AB, BA, AC, CA, BC and CB. **COMBINA** implements the formula: **(Count1+Count2-1)! / (Count2!(Count1-1)!)**