(i) Adsorption

Phage (T_4) particles interact with cell wall of host (*E. coli*). The phage tail makes contact between the two, and tail fibres recognize the specific receptor sites present on bacterial cell surface. The lipopolysaccharides of tail fibres act as receptor in phages. The process involving the recognition of phage to bacterium is called **landing**. Once the contact is established between tail fibres and bacterial cell, tail fibres bend to anchor the pins and base plate to the cell surface. This step is called **pinning.**

(ii) Penetration

The penetration process involves mechanical and enzymatic digestion of the cell wall of the host. At the recognition site phage digests certain cell wall structure by viral enzyme (lysozyme). After pinning the tail sheath contracts (using ATP) and appears shorter and thicker. After contraction of the base plate enlarges through which DNA is injected into the cell wall without using metabolic energy. The step involving injection of DNA particle alone into the bacterial cell is called **Transfection**. The empty protein coat leaving outside the cell is known as **'ghost'**.

(iii) Synthesis

This step involves the degradation of bacterial chromosome, protein synthesis and DNA replication. The phage nucleic acid takes over the host biosynthetic machinery. Host DNA gets inactivated and breaks down. Phage DNA suppresses the synthesis of bacterial protein and directs the metabolism of the cell to synthesis the proteins of the phage particles and simultaneously replication of Phage DNA also takes place.

(iv) Assembly and Maturation

The DNA of the phage and protein coat are synthesized separately and are assembled to form phage particles. The process of assembling the phage particles is known as **maturation**. After 20 minutes of infection, about 300 new phages are assembled.

(v) Release

The phage particle gets accumulated inside the host cell and are released by the lysis of host cell wall.




Figure 1.5: Multiplication cycle of phage

b. Lysogenic Cycle

In the lysogenic cycle the phage DNA gets integrated into host DNA and gets multiplied along with nucleic acid of the host. No independent viral particle is formed (Figure 1.5b).

As soon as the phage injects its linear DNA into the host cell, it becomes circular and integrates into the bacterial chromosome by recombination. The integrated phage DNA is now called **prophage.** The activity of the prophage gene is repressed by two repressor proteins which are synthesized by phage genes. This checks the synthesis of new phages within the host cell. However, each time the bacterial



Viruses infecting blue green algae are called **Cyanophages** and are first reported by Safferman and Morris in the year 1963(Example LPP1 - *Lyngbya, Plectonema* and *Phormidium*). Similarly, Hollings(1962) reported viruses infecting cultivated Mushrooms and causing die back disease. The viruses attacking fungi are called **Mycoviruses** or **Mycophages.**

XIth_BIO-BOTANY_Ch_1-EM.indd 6 5/17/2020 11:09:22 PM