protoplasm which is in gel form can change into sol form by **solation** and the sol can change into gel by **gelation**. These gel-sol conditions of colloidal system are prime basis for mechanical behaviour of cytoplasm.

1. Protoplasm is translucent, odourless and polyphasic fluid.
2. It is a crystal colloid solution which is a mixture of chemical substances forming crystalloid i.e. true solution (sugars, salts, acids, bases) and others forming colloidal solution (Proteins and lipids)
3. It is the most important property of the protoplasm by which it exhibits three main phenomena namely Brownian movement, amoeboid movement and cytoplasmic streaming or cyclosis. Viscosity of protoplasm is 2–20 centipoises. The Refractive index of the protoplasm is 1.4.
4. The pH of the protoplasm is around 6.8, contain 90% water (10% in dormant seeds)
5. Approximately 34 elements are present in protoplasm but only 13 elements are main or universal elements i.e. C, H, O, N, Cl, Ca, P, Na, K, S, Mg, I and Fe. Carbon, Hydrogen, Oxygen and Nitrogen form the 96% of protoplasm.
6. Protoplasm is neither a good nor a bad conductor of electricity. It forms a delimiting membrane in contact with water and solidifies when heated.
7. **Cohesiveness:** Particles or molecules of protoplasm are adhered with each other by forces, such as **Van der Waal's bonds**, that hold long chains of molecules together. This property varies with the strength of these forces.
8. **Contractility:** The contractility of protoplasm is important for the absorption and removal of water especially for stomatal operations.
9. **Surface tension:** The proteins and lipids of the protoplasm have less surface tension, hence they are found at the surface forming the membrane. On the other hand the chemical substances (NaCl) have high surface tension, so they occur in deeper parts of the protoplasm.

6.3.3 Cell sizes and shapes

Cell greatly vary in size, shape and also in function. Group of cells with similar structures are called **tissue** they integrate together to perform similar function, group of tissue join together to perform similar function called **organ**, group of organs with related function called **organ system**, organ system coordinating together to form an **organism**.

Shape

The shape of cell vary greatly from organism to organism and within the organism itself. In bacteria, cell shape vary from round (**cocci**) to rectangular (**rod**). In virus, shape of the envelope varies from round to hexagonal or '**T**' shaped. In fungi, globular to elongated cylindrical cells and the spores of fungi vary



1 cm = 1/100 metre
1 mm = 1/1000 metre =1/10 cm
1 µm = 1/1000,000 metre = 1/10,000 cm
1 nm = 1/1,000,000,000 metre = 1/10,000,000 cm
1 A^{o} = 1/10,000,000,000 metre =1/100,000,000 cm

or

$1\ m = 10^{2} cm = 10^{3} mm = 10^{6} \mu m = 10^{9} nm = 10^{10}\ A^{o}$

m = metre cm = centimetre mm = millimetre µm = micrometre
nm = nanometre A^{o} =Angstrom

XIth_BIO-BOTANY_Ch_6-EM.indd 122 5/17/2020 11:20:16 PM