Lampbrush chromosomes occur at the diplotene stage of first meiotic prophase in oocytes of an animal **Salamandar** and in giant nucleus of the unicellular alga *Acetabularia*. It was first observed by **Flemming** in 1882. The highly condensed chromosome forms the chromosomal axis, from which lateral loops of DNA extend as a result of intense RNA synthesis.




Figure 6.26: Lampbrush chromosomes

6.8. Flagella

6.8.1 Prokaryotic Flagellum

Bacterial flagella are helical appendages helps in motility. They are much thinner than flagella or cilia of eukaryotes. The filament contains a protein called **flagellin**. The structure consists of a basal body associated with cytoplasmic membrane and cell wall with short hook and helical filament. Bacteria rotates their helical flagella and propels rings present in the basal body which are involved in the rotary motor that spins the flagellum.

Check your grasp ?
When E.coli are cultured in medium rich in glucose they lack flagella. When grown in nutritionally poor medium they possess flagella. What does this indicate about the value of flagella?

Flagella is essential to seek out a nutritionally more favourable environment.

Structure of flagella in Bacteria

The gram positive bacteria contain only two basal rings. S-ring is attached to the inside of peptidoglycan and M-ring is attached to the cell membrane. In Gram negative bacteria two pairs of rings proximal and distal ring are connected by a central rod. They are L-Lipopolysaccharide ring, P-Peptidoglycan ring, S-Super membrane ring and M-membrane ring. The outer pair L and P rings is attached to cell wall and the inner pair S and M rings attached to cell membrane (Figure 6.27).




Figure 6.27: Structure of Bacterial Flagellum

Mechanism of flagellar movement – proton motive force

In flagellar rotation only proton movements are involved and not ATP. Protons flowing back into the cell through the basal body rings of each flagellum drives it to rotate. These rings constitute the rotary motor.The proton motive force (The force derived from the electrical potential and the hydrogen ion gradient across the cytoplasmic membrane) drives the flagellar motor. For the rotation of flagellum the energy is derived from proton gradient across the plasma membrane generated by oxidative phosphorylation. In bacteria flagellar motor is located in the plasma membrane where the oxidative phosphorylation takes place.Therefore, plasma membrane is a site of generation of proton motive force.

6.8.2 Eukaryotic Flagellum– Cell Motility

Structure

Eukaryotic Flagella are enclosed by unit membrane and it arises from a basal body. Flagella is composed of outer nine pairs of microtubules with two microtubules in its centre (9+2 arrangement). Flagella are microtubule projection of the plasma membrane. Flagellum is longer than cilium (as long as 200μm). The structure of flagellum

XIth_BIO-BOTANY_Ch_6-EM.indd 136 5/17/2020 11:20:37 PM