


Test for starch

a. Test on potato; b. test on starch at varied concentrations; c. starch – iodine reaction

8.3.6 Celluloses

Cellulose is a structural polysaccharide made up of thousands of glucose units. In this case, β-glucose units are held together by 1,4 glycosidic linkage, forming long unbranched chains. Cellulose fibres are straight and uncoiled. It has many industrial uses which include cellulose fibres as cotton, nitrocellulose for explosives, cellulose acetate for fibres of multiple uses and cellophane for packing (Figure 8.7).




Figure 8.7: Cellulose molecule

8.3.7 Chitin

Chitin is a homo polysaccharide with amino acids added to form **mucopolysaccharide**. The basic unit is a nitrogen containing glucose derivative known as **N-acetyl glucosamine.** It forms the exoskeleton of insects and other arthropods. It is also present in the cell walls of fungi (Figure 8.8).





Figure 8.8: Structure of Chitin molecule



8.3.8 Test for Reducing Sugars

Aldoses and ketoses are reducing sugars. This means that, when heated with an alkaline solution of copper (II) sulphate (a blue solution called **benedict's solution**), the aldehyde or ketone group reduces Cu^{2+} ions to Cu^{+} ions forming brick red precipitate of copper(I) oxide. In the process, the aldehyde or ketone group is oxidised to a carboxyl group (–COOH). This reaction is used as test for reducing sugar and is known as **Benedict's test.** The results of benedict's test depends on concentration of the sugar. If there is no reducing sugar it remains blue.

- Sucrose is not a reducing sugar
- The greater the concentration of reducing sugar, the more is the precipitate formed and greater is the colour change.

XIth_BIO-BOTANY_Ch_8-EM.indd 154 5/17/2020 11:22:19 PM