but each flagellum in bacteria is made up of a single fibril. Flagella are used for locomotion. Based on the number and position of flagella there are different types of bacteria (Figure 1.8)

Fimbriae or Pili

Pili or fimbriae are hair like appendages found on surface of cell wall of gram-negative bacteria (Example: *Enterobacterium*). The pili are 0.2 to 20 µm long with a diameter of about 0.025µm. In addition to normal pili there are special type of pili which help in conjugation called sex pili are also found.

1.4.4 Gram staining procedure

The Gram staining method to differentiate bacteria was developed by Danish Physician Christian Gram in the year1884. It is a differential staining procedure and it classifies bacteria into two classes - Gram positive and Gram negative. The steps involved in Gram staining procedure is given in Figure 1.10. The Gram positive bacteria retain crystal violet and appear dark violet whereas Gram negative type loose the crystal violet and when counterstained by safranin appear red under a microscope.

Most of the gram positive cell wall contain considerable amount of teichoic acid and teichuronic acid. In addition, they may contain



Figure 1.10: Steps involved in Gram Staining

polysaccharide molecules. The gram negative cell wall contains three components that lie outside the peptidoglycan layer. 1. Lipoprotein 2. Outer membrane 3.Lipopolysaccharide. Thus the different results in the gram stain

Table 1.6: Difference between Gram Positive and Gram Negative Bacteria

S. No.	Characteristics	Gram positive Bacteria	Gram negative Bacteria
1.	Cell wall	Thick layered with (0.015 µm-0.02µm)	Thin layered with (0.0075µm–0.012µm)
2.	Rigidity of cell wall	Rigid due to presence of Peptidoglycans	Elastic due to presence of lipoprotein-polysaccharide mixture
3.	Chemical composition	Peptidoglycans-80% Polysaccharide-20% Teichoic acid present	Peptidoglycans-3 to 12% rest is polysaccharides and lipoproteins. Teichoic acid absent
4.	Outer membrane	Absent	Present
5.	Periplasmic space	Absent	Present
6.	Susceptibility to penicillin	Highly susceptible	Low susceptible
7.	Nutritional requirements	Relatively complex	Relatively simple
8.	Flagella	Contain 2 basal body rings	Contain 4 basal body rings
9.	Lipid and lipoproteins	Low	High
10.	Lipopolysaccharides	Absent	Present

XIth_BIO-BOTANY_Ch_1-EM.indd 13 5/17/2020 11:09:31 PM