


Figure 14.3: Molecular structure of ATP

14.3 Redox Reactions

$NAD^+ + 2e^- + 2H^+ \longrightarrow NADH + H^+$

$FAD + 2e^- + 2H^+ \longrightarrow FADH_2$

When NAD^+ (Nicotinamide Adenine Dinucleotide-oxidised form) and FAD (Flavin Adenine Dinucleotide) pick up electrons and one or two hydrogen ions (protons), they get reduced to $NADH + H^+$ and $FADH_2$ respectively. When they drop electrons and hydrogen off they go back to their original form. The reaction in which NAD^+ and FAD gain (reduction) or lose (oxidation) electrons are called **redox reaction** (Oxidation reduction reaction). These reactions are important in cellular respiration.

14.4 Types of Respiration

Respiration is classified into two types as aerobic and anaerobic respiration (Figure 14.4)

Handy mnemonic



LEO the lion says GER

LEO - Loss of Electrons is Oxidation

GER - Gain of Electrons is Reduction




Figure 14.4: Types of Respiration

14.4.1 Aerobic respiration

Respiration occurring in the presence of oxygen is called **aerobic respiration**. During aerobic respiration, food materials like carbohydrates, fats and proteins are completely oxidised into CO_2, H_2O and energy is released. Aerobic respiration is a very complex process and is completed in four major steps:

1. Glycolysis
2. Pyruvate oxidation (Link reaction)
3. Krebs cycle (TCA cycle)
4. Electron Transport Chain (Terminal oxidation).

14.4.2 Anaerobic respiration

In the absence of molecular oxygen glucose is incompletely degraded into either ethyl alcohol or lactic acid (Table 14.1). It includes two steps:

1. Glycolysis 2. Fermentation

Table 14.1: Differences between aerobic and anaerobic respiration

Aerobic respiration	Anaerobic Respiration
1. It occurs in all living cells of higher organisms.	It occurs yeast and some bacteria.
2. It requires oxygen for breaking the respiratory substrate.	Oxygen is not required for breaking the respiratory substrate.
3. The end products are CO_2 and H_2O.	The end products are alcohol, and CO_2 (or) lactic acid.
4. Oxidation of one molecule of glucose produces 36 ATP molecules.	Only 2 ATP molecules are produced.
5. It consists of four stages-glycolysis, link reaction, TCA cycle and electron transport chain.	It consists of two stages-glycolysis and fermentation.
6. It occurs in cytoplasm and mitochondria.	It occurs only in cytoplasm.

XIth_BIO-BOTANY_Ch_14-EM.indd 263 5/17/2020 11:29:34 PM