
Therefore, in plant cells net yield of 30 ATP molecules for complete aerobic oxidation of one molecule of glucose. But in those animal cells (showing malate shuttle mechanism) net yield will be 32 ATP molecules.



Cyanide resistant respiration is believed to be responsible for the climacteric in fruits

Cyanide resistant respiration is known to generate heat in thermogenic tissues.

The amount of heat produced in thermogenic tissues may be as high as 51°C.



Peter Mitchel, a British Biochemist received Nobel prize for Chemistry in 1978 for his work on the coupling of oxidation and phosphorylation in mitochondria.

14.6 Respiratory Quotient (RQ)

The ratio of volume of carbon dioxide given out and volume of oxygen taken in during respiration is called **Respiratory Quotient or Respiratory ratio**. RQ value depends upon respiratory substrates and their oxidation.

$$RQ = \frac{\text{Volume of } CO_2 \text{ liberated}}{\text{Volume of } O_2 \text{ consumed}}$$

1. The respiratory substrate is a carbohydrate, it will be completely oxidised in aerobic respiration and the value of the RQ will be equal to unity.

$C_6H_{12}O_6 + 6O_2 \longrightarrow 6CO_2\uparrow + 6H_2O + \text{Energy}$

Glucose

$$\text{RQ of glucose} = \frac{6 \text{ molecules of } CO_2}{6 \text{ molecules of } O_2}$$

$$= 1 \text{ (unity)}$$

2. If the respiratory substrate is a carbohydrate it will be incompletely oxidised when it goes through anaerobic respiration and the RQ value will be infinity.

$C_6H_{12}O_6 \longrightarrow 2CO_2\uparrow + 2C_2H_5OH + \text{Energy}$

Glucose Ethyl alcohol

$$\left.\begin{matrix}\text{RQ of glucose} \\ \text{Anaerobically}\end{matrix}\right\} = \frac{2 \text{ molecules of } CO_2}{\text{zero molecule of } O_2}$$

$$= \infty \text{ (infinity)}$$

3. In some succulent plants like *Opuntia, Bryophyllum* carbohydrates are partially oxidised to organic acid, particularly malic acid without corresponding release of CO_2 but O_2 is consumed hence the RQ value will be zero.

$2C_6H_{12}O_6 + 3O_2 \longrightarrow 3C_4H_6O_5 + 3H_2O + \text{Energy}$

Glucose Malic acid

$$\text{RQ of glucose in succulents} = \frac{\text{zero molecule of } CO_2}{3 \text{ molecules of } O_2}$$

$$= 0 \text{ (zero)}$$

4. When respiratory substrate is protein or fat, then RQ will be less than unity.

$2(C_{51}H_{98}O_6) + 145O_2 \longrightarrow 102CO_2\uparrow + 98H_2O + \text{Energy}$

Tripalmitin(Fat)

$$\text{RQ of Tripalmitin} = \frac{102 \text{ molecules of } CO_2}{145 \text{ molecules of } O_2}$$

$$= 0.7 \text{ (less than unity)}$$

5. When respiratory substrate is an organic acid the value of RQ will be more than unity.

$C_4H_6O_5 + 3O_2 \longrightarrow 4CO_2\uparrow + 3H_2O + \text{Energy}$

Malic acid

$$\text{RQ of malic acid} = \frac{4 \text{ molecules of } CO_2}{3 \text{ molecules of } O_2}$$

$$= 1.33 \text{ (more than unity)}$$

Significance of RQ

1. RQ value indicates which type of respiration occurs in living cells, either aerobic or anaerobic.
2. It also helps to know which type of respiratory substrate is involved.



Red colour in various parts of plants is due to the presence of anthocyanin, synthesis of which require more O_2 than CO_2 evolved. RQ will be less than one.

XIth_BIO-BOTANY_Ch_14-EM.indd 270 5/17/2020 11:29:37 PM