| | |
|---|---|
| **Protologue** | Set of information associated with the scientific name of a taxon at its first valid publication containing the entire original material regarding the taxon |
| **Quantasome** | Morphological expression of physiological photosynthetic units, located on the inner membrane of thylakoid lamellae. Act as photosynthetic unit contains 200 to 300 chlorophyll molecules. |
| **Quantum** | The energy contained in a photon is represented as quantum |
| **Quantum requirement** | The number of photons or quanta required to release one molecule of oxygen during photosynthesis |
| **Quantum yield** | The number of oxygen molecules produced per quantum of light absorbed. |
| **Quiescent centre concept** | Inactive region of root meristem |
| **Rachilla** | Central axis of a spikelet |
| **Radial vascular bundles** | Xylem and phloem present on different radii |
| **Ray Parenchyma** | Parenchyma cells arranged in radial rows |
| **Redox reactions** | Oxidation (loss of electrons) and Reduction (gain of electrons) reactions are called redox reactions. |
| **Reduction** | CO2 is reduced into Carbohydrates (gain of electrons) |
| **Rib-meristem** | Meristem which divides anticlinally in two planes |
| **RUBISCO** | Enzyme responsible for fixation of Carbon dioxide, the most abundant protein (Ribulose 1,5 bisphosphate Carboxylase Oxygenase) |
| **Salt stress** | Adverse effects of excess mineral salts on plants |
| **Sap** | It is a fluid consist of water and dissolved minerals |
| **Slime body** | A special protein (Phloem Protein) in sieve tubes |
| **Sporophyte** | Diploid plant body |
| **Stellate hairs** | Star shaped hairs |
| **Stratification** | A process of breaking the dormancy of some plants resulting from chilling requirements |
| **Subsidiary cells** | Surrounding guard cells in the leaf epidermis |
| **Sucrose** | Non-reducing disaccharide composed of glucose and fructose |
| **Teloemorph** | Sexual or perfect state of the fungi |
| **Thallospores** | Asedual spores formed due to the fragmentation of hyphae |
| **Trichoblasts** | One type of epidermal cells that is also called short cell |
| **Trichomes** | Unicellular or multicellular appendages |
| **Triplicate** | Pollen grain with three furrows or pores |
| **Tunica-carpus theory** | Two zones of apical meristem Tunica and Carpus |
| **X-Ray crystallography** | Most commonly used technique for determining the three-dimensional structure of macromolecules (particularly proteins and nucleic acids) by passing x-rays |
| **Xylos** | Wood |
| **Zoospore** | Motile, asexual spores |
| **Zygospore** | Thick walled diploid resting spores |



XIth_BIO-BOTANY_Ch_15-EM.indd 298 5/17/2020 11:30:35 PM