on Bark) **Lignicolous**(on Wood) **Saxicolous**(on rocks) **Terricolous**(on ground) Marine(on siliceous rocks of sea) Fresh water(on siliceous rock of fresh water).

- On the basis of morphology of the thallus they are divided into **Leprose** (a distinct fungal layer is absent) **Crustose**-crust like; Foliose-leaf like; **Fruticose**- branched pendulous shrub like (Figure 1.25).



(a) Crustose lichen

(b) Foliose Lichen

(c) Fruticose Lichen

Figure 1.25: Types of Lichens

- The distribution of algal cells distinguishes lichens into two forms namely **Homoiomerous** (Algal cells evenly distributed in the thallus) and **Heteromerous** (a distinct layer of algae and fungi present).
- If the fungal partner of lichen belongs to ascomycetes, it is called **Ascolichen** and if it is basidiomycetes it is called **Basidiolichen**.

Lichens secrete organic acids like Oxalic acids which corrodes the rock surface and helps in weathering of rocks, thus acting as pioneers in Xerosere. Usnic acid produced from lichens show antibiotic properties. Lichens are sensitive to air pollutants especially to sulphur-di-oxide. Therefore, they are considered as pollution indicators. The dye present in litmus paper used as acid base indicator in the laboratories is obtained from *Roccella montagnei. Cladonia rangiferina* (Reindeer mose) is used as food for animals living in Tundra regions.

Summary

Earth is endowed with living and nonliving things. The attributes of living things include growth, metabolism, reproduction, irritability and so on. Viruses are considered as Biological puzzle and exhibit both living and non living characteristic features. They are ultramicroscopic, obligate parasites and cause disease in plants and animals. They multiply by lytic and lysogenic cycle.

Five Kingdom classification was proposed by Whittaker, which include Monera, Protista, Fungi, Plantae and Animalia. Carl woese divided the living world into 3 domains- Bacteria, Archaeae and Eukarya. The domain Eukarya include Plantae, Animalia and Fungi. A new Kingdom called Chromista was erected to include Diatoms, Cryptomonads and Oomycetes. Bacteria are microscopic, prokaryotic organisms and possess peptidoglycan in their cell wall. Based on Gram Staining method they are classified into Gram positive and Gram negative type. They reproduce asexually by binary fission. Sexual reproduction occurs through conjugation, transformation and transduction. Archaebacteria are prokaryotic and are adapted to thrive in extreme environments.

Cyanobacteria are prokaryotic organisms and are also called Blue Green Algae. The members of this group are ensheathed by mucilage cover. They reproduce by vegetative and asexual methods.

Fungi are eukaryotic, heterotrophic, unicellular or multicellular organisms. The cell wall is made up of chitin. They reproduce asexually by producing sporangiospores, conidia, thallospores, chlamydospores etc., The sexual reproduction is isogamous, ansiogamous and oogamous. In addition, gametic copulation, gametic fusion, spermatisation are also found. They are

XIth_BIO-BOTANY_Ch_1-EM.indd 28 5/17/2020 11:09:47 PM