


internodes only. Their stem nature is evident by the fact that they bear buds, scales and flowers. Example: *Asparagus* (cylindrical cladode), *Ruscus* (flattened cladode).

6. Thorns

Thorn is a woody and sharp pointed modified stem. Either the axillary bud or the terminal bud gets modified into thorns. In *Citrus* and *Atalantia* axillary bud is modified into thorns.

II. Sub aerial stem modifications

Sub aerial stem found in plants with weak stem in which branches lie horizontally on the ground. These are meant for vegetative propagation. They may be sub aerial or partially sub terranean.

1. Runner

This is a slender, prostrate branch creeping on the ground and rooting at the nodes. Example: *Oxalis* (Wood sorrel), lawn grass (*Cynodon dactylon*).

2. Stolon

This is also a slender, lateral branch originating from the base of the stem. But it first grows obliquely above the ground, produces a loop and bends down towards the ground. When touches the ground it produces roots and becomes an independent plantlet. Example: *Mentha piperita* (peppermint), *Fragaria indica* (wild strawberry).




Figure 3.9: (a) Runner-*Oxalis* **(b)** Stolon-*Fragaria*

3. Sucker

Sucker develops from an underground stem and grows obliquely upwards and gives rise to a separate plantlet or new plant. Example: *Chrysanthemum, Bambusa.*

4. Offset

Offset is similar to runner but found in aquatic plants especially in rosette leaved forms. A short thick lateral branch arises from the lower axil and grows horizontally leafless for a short distance, then it produces a bunch of rosette

XIth_BIO-BOTANY_Ch_3-EM.indd 55 5/17/2020 11:13:00 PM