
Leaf hooks-*Bignonia*


Leaf spines- *Opuntia*


Phyllode-*Acacia*


Pitcher-*Nepenthes*

Figure 3.17: Leaf Modification

VI. Pitcher

The leaf becomes modified into a pitcher in *Nepenthes* and *Sarracenia*. In *Nepenthes* the basal part of the leaf is laminar and the midrib continues as a coiled tendrillar structure. The apical part of the leaf is modified into a pitcher the mouth of the pitcher is closed by a lid which is the modification of leaf apex.

VII. Bladder

In bladderwort (*Utricularia*), a rootless free-floating or slightly submerged plant common in many water bodies, the leaf is very much segmented. Some of these segments are modified to form bladder-like structures, with a trap-door entrance that traps aquatic animalcules.

VIII Floral leaves

Floral parts such as sepals, petals, stamens and carpels are modified leaves. Sepals and petals are leafy. They are protective in function and considered non-essential reproductive parts. Petals are usually coloured which attract the insects for pollination. Stamens are considered pollen bearing microsporophylls and carpels are ovule bearing megasporophylls.

3.7.7 Leaf duration

Leaves may stay and function for few days to many years, largely determined by the adaptations to climatic conditions.

Caducuous (Fagacious)

Falling off soon after formation. Example: *Opuntia, Cissus quadrangularis.*

Deciduous

Falling at the end of growing season so that the plant (tree or shrub) is leafless in winter/ summer season. Example: *Maple, Plumeria, Launea, Erythrina.*

Evergreen

Leaves persist throughout the year, falling regularly so that tree is never leafless. Example: *Mimusops, Calophyllum.*

Marcescent

Leaves not falling but withering on the plant as in several members of Fagaceae.

Summary

Flowering plants consist of two major organ systems: Underground root system and aerial shoot system. Roots perform the functions of anchoring and absorbing nutrients from the soil. However some roots perform additional functions for which they undergo various modifications in shape, form and structure. Tap root continue the growth from the radicle which further branches into secondary roots. Adventitious roots arise from different parts of the plant other than radicle. Stem helps to display the leaves to get maximum sunlight and positioning flowers and fruits to attract pollination and dispersal agents. Apart from the normal functions the stems are modified to perform various functions such as food storage, perennation and protection. Leaves are exogenous in origin and function as food synthesizing and gaseous exchange sites. Some leaves also perform additional functions for which they are modified in their morphology.

XIth_BIO-BOTANY_Ch_3-EM.indd 62 5/17/2020 11:13:16 PM