Hypogynous: The term is used for sepals, petals and stamens attached at the base of a superior ovary. Example: Malvaceae	**Epihypogynous:** The term is used for sepals, petals and stamens attached at the middle of the ovary (half-inferior). Example: Fabaceae, Rosaceae.	**Epigynous**: The term is used for sepals, petals and stamens attached at the tip of an inferior ovary. Example: Cucumber, Apple, Asteraceae.
Perigynous: The term is used for a hypanthium attached at the base of a superior ovary.	**Epiperigynous:** The term is used for hypanthodium attached at the apex of an inferior ovary.	

Placentation

The mode of distribution of placenta inside the ovary



Marginal
It is with the placentae along the margin of a unicarpellate ovary.
Example: *Fabaceae.*



Superficial
Ovules arise from the surface of the septa.
Example: *Nymphaeaceae.*



Free-central
It is with the placentae along the column in a compound ovary without septa.
Example: *Caryophyllaceae, Dianthus, Primerose.*



Axile
The placentae arises from the column in a compound ovary with septa.
Example: *Hibiscus, Tomato Lemon.*



Parietal
It is the placentae on the ovary walls or upon intruding partitions of a unilocular, compound Ovary.
Example: *Mustard, Argemone, Cucumber.*



Basal
It is the placenta at the base of the ovary.
Example: *Sunflower (Asteraceae) Marigold*

4.5.6 Perianth / Androecial position on thalamus:

It describes placement of the perianth and androecium relative to the ovary and to a hypanthium, if present (Figure 4.25).

Hypanthium (Staminal disk) : a fleshy elevated often nectariferous cup like thalamus

4.6 Construction of floral diagram and floral formula

A floral formula is a simple way to explain the salient features of a flower. The floral diagram is a representation of the cross section of the flower. It represents floral whorls arranged as viewed from above. Floral diagram shows the number and arrangement of bract, bracteoles and floral parts, fusion, overlapping and placentation.

The branch that bears the flower is called **mother axis**.

The side of the flower facing the mother axis is called **posterior side**. The side facing the bract is the anterior side.

The members of different floral whorls are shown arranged in concentric rings.

Br : Bracteate.

Ebr : Ebracteate

Brl : Bracteolate

Ebrl : Ebracteolate

⊕ : Actinomorphic

% : Zygomorphic

XIth_BIO-BOTANY_Ch_4-EM.indd 77 5/17/2020 11:15:34 PM