


Figure 4.27: Classification of fruits based on formation

epicarp, mesocarp and endocarp. It is subdivided into the following.


a) Berry: Fruit develops from bicarpellary or multicarpellary, syncarpous ovary. Here the epicarp is thin, the mesocarp and endocarp remain undifferentiated. They form a pulp in which the seeds are embedded. Example: Tomato, Grapes, Brinjal.

b) Drupe: Fruit develops from monocarpellary, superior ovary. It is usually one seeded. Pericarp is differentiated into outer skinny epicarp, fleshy and pulpy mesocarp and hard and stony endocarp around the seed. Example: Mango, Coconut.

c) Pepo: Fruit develops from tricarpellary inferior ovary. Pericarp turns leathery or woody which encloses, fleshy mesocarp and smooth endocarp. Example: Cucumber, Watermelon, Bottle gourd, Pumpkin.

d) Hesperidium: Fruit develops from multicarpellary, multilocular, syncarpous, superior ovary. The fruit wall is differentiated into leathery epicarp with oil glands, a middle fibrous mesocarp. The endocarp forms distinct chambers, containing juicy hairs. Example: Orange, Lemon.

e) Pome: It develops from multicarpellary, syncarpous, inferior ovary. The receptacle



Berry (Tomato) **Drupe** (Mango) **Pepo** (Cucumber)

Hesperidium (Orange) **Pome** (Apple) **Balausta** (Pomegranate)

Figure 4.28: Simple fleshy fruits

XIth_BIO-BOTANY_Ch_4-EM.indd 79 5/17/2020 11:15:35 PM