also develops along with the ovary and becomes fleshy, enclosing the true fruit. In pome the epicarp is thin skin like and endocarp is cartilagenous. Example: Apple, Pear.

f) Balausta: A fleshy indehiscent fruit developing from multicarpellary, multilocular inferior ovary whose pericarp is tough and leathery. Seeds are attached irregularly with testa being the edible portion. Example: Pomegranate.

B. Dry Fruit

They develop from single ovary where the pericarp is dry and not differentiated into epicarp, mesocarp and endocarp. It is further subdivided into three types.

1) Dry dehiscent fruit

Pericarp is dry and splits open along the sutures to liberate seeds. They can be classified into following types.

a) Follicle: Fruit develops from mono-carpellary, superior ovary and dehisces along one suture. Example: *Calotropis.*


b) Legume or pod: Fruit develops from monocarpellary, superior ovary and dehisces through both dorsal and ventral sutures. Example: *Pisum.*

c) Siliqua: Fruit develops from bicarpellary, syncarpous, superior ovary initially one chambered but subsequently becomes two chambered due to the formation of false septum (**replum**). The fruit dehisces along two suture. Example: *Brassica.*

d) Silicula: Fruit similar to siliqua but shorter and broader. Example: *Capsella.*

e) Capsule: Fruit develops from multicarpellary, syncarpous, superior ovary. Based on the dehiscence pattern they are divided into.

i) Septicidal: Capsule splitting along septa and valves remaining attached to septa. Example: *Aristolochia.*



Follicle (*Calotropis*) **Legume** (*Pisum*)
Siliqua (*Brassica*) **Silicula** (*Capsella*)
Loculicidal (*Abelmoschus*) **Septifragal** (*Datura*)

Figure 4.29: Dry dehiscent fruit

ii) Loculicidal: Capsule splitting along locules and valves remaining attached to septa. Example: *Abelmoschus*.

iii) Poricidal: Dehiscence through terminal pores. Example: *Papaver*.

2) Dry indehiscent fruit

Dry fruit which does not split open at maturity. It is subdivided into.

a) Achene: Single seeded dry fruit developing from single carpel with superior ovary. Achenes commonly develop from apocarpous pistil, Fruit wall is free from seed coat. Example: *Clematis, Delphinium*.

b) Cypsela: Single seeded dry fruit, develops from bicarpellary, syncarpous, inferior ovary with reduced scales, hairy or feathery calyx lobes. Example: *Tridax.*

c) Caryopsis: It is a one seeded fruit which develops from a monocarpellary,

XIth_BIO-BOTANY_Ch_4-EM.indd 80 5/17/2020 11:15:36 PM