Table 8.1 Comparison of fat and and water soluble vitamins

S. No.	Characteristics	Fat soluble vitamins	Water soluble vitamins
1	Solubility	Fat soluble	Water soluble
2	Absorption	Bile salts are required	Simple intestinal absorption
3	Transportation	Transported by carrier protein	Travels freely in the body without requiring carrier protein (except Vitamin B_{12})
4	Storage	Stored in liver and fatty tissues	Not stored (except vitamin B_{12})
5	Excretion	Usually the surplus vitamins are stored	Surplus vitamins are detected in kidney and removed in urine
6	Accumulation	Usually hypervitaminosis occurs	Usually hypervitaminosis deosn't occur (except in high dosage and slow release of some B vitamins)
7	Deficiency compensation	Required in periodic doses (weeks or months)	Required in frequent doses (1 - 3 days)

Sources:

Liver, oil, butter, milk, egg-yolk, tomato, carrot, green yellow vegetables, spinach and fruits such as mango and papaya (Figure 8.2).

Meat : 25000 IU / 100 g

Spinach (Cooked): 1200 IU / 100 g

Beta Carotene
Carrots (Cooked) : 2500 IU / 100 g

Drumstick leaves: 7000-8000 microgram / 100g

Unit 8.indd 198 21-05-2018 15:11:36