




Golden langur

Trachypithecus geei

The golden langur's coat varies from cream-colored in summer to burnished gold in winter. This elusive, long-tailed monkey was not recognized as a species until the 1950s, and very little is known about it even today. Golden langurs live in groups of 3–40, and rarely come to the ground, a strategy that helps them avoid predators such as tigers. They are severely threatened by habitat loss.

- ↔ 19–28 in (49–72 cm)
- ⚖ 21–26 lb (9.5–12 kg)
- ⊗ Endangered
- 🌿 Leaves, buds, fruit, seeds
- 🏠



S. Asia

◁ TREETOP DWELLER

Golden langurs spend most of their time high in the forest canopy, only rarely descending to the ground to drink or lick up mineral salts.




LOCATION



CLIMATE

The region is temperate, with cool summers and colder winters. Rain falls mostly during summer monsoons; winter snows prevail at higher levels.



Bhutan takin

Budorcas whitei

barrel-shaped body covered in shaggy hair



During spring, large mixed herds of takin—robust relatives of wild goats—congregate in sunny clearings high up in bamboo forests. As winter approaches, they fragment into fours and fives and head for lower areas. If threatened, they retreat into dense bamboo thickets and lie down.

- ↔ 6–7 ft (1.8–2.1m)
- ⚖ 330–772 lb (150–350 kg)
- ⊗ Vulnerable
- 🌿 Forbs, shrubs, trees
- 🏠



S. Asia

both sexes have short, thick horns

▷ ATYPICAL BODY

The takin's body shape is said to be primitive compared to most hooved species; it has short, stocky legs and a rounded snout.



bird and 300 mammal species

CHANGING ALTITUDES
Many species migrate up and down the Himalayas to avoid the worst of the winter weather and exploit summer food sources. One example is the satyr tragopan, a pheasant that moves to low-altitude forests in winter and high areas in summer.

SATYR TRAGOPAN

