





△ GOING FISHING
These stocky little penguins use their short, strong legs to make double-footed kangaroo hops across rocks when going to and from the ocean.



Imperial shag

Phalacrocorax atriceps



The imperial shag nests on rocky headlands and islands and mostly feeds quite close to the shore. Like other shags and cormorants, it has relatively heavy bones and little body fat—this reduces

buoyancy and makes it a more efficient underwater forager. Unlike terns and gannets, which locate fish from the air by sight, the shag dives deep and searches systematically for prey.

↔ 27–30 in (69–76 cm)
⚖ 6–8 lb (2.7–3.6 kg)
⊗ Common
🐟 Fish
🏠 

S. South America, Southern Ocean islands, Antarctic Peninsula



△ EGG CUPS
Each nest is a mound of seaweed, grass, and mud liberally mixed with white excrement, with a shallow bowl for two or three eggs.

Antarctic tern

Sterna vittata

The southern equivalent of the Arctic tern, the Antarctic tern does not undertake vast migrations from north to south. It breeds in November and December when the northern species is

“wintering” at sea. Some birds remain close to nesting colonies, while others move far out to sea, feeding along the edges of the pack ice and often resting on ice floes.

↔ 14–16 in (36–41 cm)
⚖ 5–6 oz (150–175 g)
⊗ Common
🐟 Fish
🏠 

SE. South America, Southern Africa, Southern Ocean islands, Antarctic Peninsula



◁ BREEDING ADULT
In its summer plumage, the Antarctic tern looks very similar to its Arctic cousin. It breeds on rocky islands in the Southern Ocean.