

Myanmar snub-nosed monkey

Rhinopithecus strykeri

- ↔ 22 in (56 cm)
- 📍 S. Asia
- ⊗ Critically endangered

Discovered in 2010 in northern Myanmar, this rare and endangered monkey is mostly black, with a contrasting white mustache, beard, and ear tufts. It has an upturned nose, which, according to local people, makes it prone to sneezing in rainy weather. It lives in mountain forests in summer, and spends the winter at lower, warmer altitudes. Its known population is less than 500, making it vulnerable to extinction.

Bengal gray langur

Semnopithecus entellus

- ↔ 20–31 in (50–78 cm)
- 📍 India
- ⊗ Common

Also known as the hanuman langur, this slender, long-tailed monkey has a striking black face that contrasts with its gray or brown fur. It occurs in a wide variety of habitats ranging from semideserts to tropical forests, often living near villages, where it makes the most of any discarded food. Elsewhere, it feeds mainly on leaves and fruit. It lives in groups of varying sizes, spending most of the daylight hours on the ground, but sleeping in trees at night.



Chacma baboon

Papio ursinus

- ↔ 24–32 in (60–82 cm)
- 📍 Southern Africa
- ⊗ Common

Found in open habitats in southern Africa, this is the biggest of the baboons—ground-dwelling monkeys with long, doglike snouts and protruding nostrils that typically live in large troops. The chacma baboon mostly forages on the ground by day and has a broad diet ranging from fruit and seeds to small gazelles, although hunting is rare. Intelligent and adaptable, it may throw stones to deter intruders.

Celebes crested macaque

Macaca nigra

- ↔ 21–23 in (53–58 cm)
- 📍 SE. Asia
- ⊗ Critically endangered

Covered with black fur but having big, expressive, red-brown eyes, this short-tailed monkey lives in tropical forests. A crest runs from the forehead back over the crown. Usually flat, the crest rises when the animal is aroused. A sociable animal, it lives in groups of 60–80, but sometimes forms mixed-sex troops of 100 or more. It mainly feeds on fruit and invertebrates, but it may also eat other small animals.



Guereza

Colobus guereza

- ↔ 21–23 in (53–58 cm)
- 📍 C. and E. Africa
- ⊗ Locally common

Mainly black with a white ruff around its naked black face, this monkey is also known as the eastern black and white colobus. It has a fringe of long, white, silky hair along its flanks and a matching white tuft at the tip of its long tail. It mainly eats tough, fibrous leaves, which it can digest thanks to a complex, three-part stomach containing gut microbes that break down the fiber.

Lar gibbon

Hylobates lar

- ↔ 17–23 in (42–59 cm)
- 📍 SE. Asia
- ⊗ Endangered

Like all gibbons, this is a superb arboreal acrobat, able to swing through trees using its long arms and, thanks to its opposable big toes, walk upright along branches. It usually lives in monogamous pairs, which reinforce their pair-bond each morning with loud hooting duets that are repeated many times. Deforestation and hunting by humans are major threats to this primate.

Sumatran orangutan

Pongo abelii

- ↔ 4–5 ft (1.2–1.5 m)
- 📍 N. Sumatra
- ⊗ Critically endangered

Slimmer than the Bornean orangutan, and with longer chestnut hair, the Sumatran orangutan is also more sociable than its largely solitary relative. Big groups may come together to feed in fruiting fig trees, moving slowly through the canopy and often using their weight to bend a branch within reach of the next one. These orangutans are almost exclusively arboreal—females virtually never travel on the

ground and adult males rarely do so. Now restricted to the north of their native island, Sumatran orangutans are critically endangered by the destruction of their rainforest habitat to create oil palm plantations.

grasping hands

coarse, shaggy coat

