

## Sumatran rhinoceros

*Dicerorhinus sumatrensis*

- ↔ 8–10 ft (2.4–3 m)
- 📍 S. and SE. Asia
- ⊗ Critically endangered

The smallest rhinoceros, and the hairiest, this two-horned species was once widespread across Southeast Asia, but is now extremely rare and localized. It is a solitary animal of forests and swamps, spending the day in a mud wallow to keep cool. It feeds mainly in the evening and early morning, gathering leaves, tender shoots, and fruit, and often uses its weight to push over young trees so it can get at their foliage.

## Lowland tapir

*Tapirus terrestris*

- ↔ 6–7 ft (1.8–2.1 m)
- 📍 N. and C. South America
- ⊗ Vulnerable



This bristly, brown-coated animal has white-tipped ears and a short, narrow mane. Like other tapirs, it has a long, mobile snout, which it uses to browse selectively for nutritious leaves, shoots, and fruit. It prefers waterside habitats and is a good swimmer—it dives into the water to escape predators such as jaguars, but is often taken by the black caiman. Females give birth to single young, which are born with white spots and stripes. These provide camouflage in the dappled shade.

## Malayan tapir

*Tapirus indicus*

- ↔ 6–8 ft (1.8–2.4 m)
- 📍 SE. Asia
- ⊗ Endangered

The largest and only Old World tapir, this species is black with a sharply contrasting white back and rump. The effect is striking, but in the Malayan tapir's shady forest habitat, it acts as camouflage, breaking up the animal's outline so it is not recognizable by predators such as tigers. The Malayan tapir is a solitary browser that feeds on a variety of soft twigs and young leaves of bushes and saplings, as well as fallen fruit.

## Giant forest hog

*Hylochoerus meinertzhageni*

- ↔ 4–7 ft (1.2–2.1 m)
- 📍 W., C., and E. Africa
- ⊗ Locally common

Perfectly described by its name, the giant forest hog is the biggest of the wild pigs and lives almost entirely in the African tropical forests. Its dark skin is covered with coarse black hair, and it has enlarged canine teeth forming tusks that grow up and out from each jaw. Unlike most pigs, it does not root in the ground for food, but eats grasses, leaves, and occasionally cultivated crops.

## Bushpig

*Potamochoerus larvatus*

- ↔ 4–5 ft (1.2–1.5 m)
- 📍 E., C., and Southern Africa
- ⊗ Common

The long-snouted bushpig is similar to the Eurasian wild boar, with a coat of coarse dark hair, a paler, bristly mane, and a pale head with white face markings. It lives in forests and swamps, in groups of up to 12 that usually consist of several females, juveniles, and a single adult male. Bushpigs use their snouts to poke around in the soil for roots, bulbs, tubers, and insect grubs, but may also scavenge for carrion.

## Moluccan babirusa

*Babyrusa babyrussa*

- ↔ 3–4 ft (0.9–1.2 m)
- 📍 SE. Asia
- ⊗ Vulnerable

This wild pig is notable for the male's tusks, which curve upward from its upper jaw to its forehead; they may be up to 12 in (30 cm) long. It has shorter, sharper tusks in its lower jaw for fighting rivals, and uses the upper tusks for defense. Males usually live alone, but females and their young travel in small groups. They eat a range of leaf, root, fruit, and animal foods, but do not root in the ground for them.

## Collared peccary

*Pecari tajacu*

- ↔ 30–39 in (75–100 cm)
- 📍 SW. US to S. South America
- ⊗ Common

Built like a small wild boar, with a barrel-shaped body, slim legs, and a pale band around its neck, this omnivore thrives in a range of habitats. Extremely sociable, it lives in tightly knit mixed-sex groups for mutual defense against enemies such as pumas. It feeds mostly on plant material, but also eats worms, lizards, and snakes.

## Pygmy hippopotamus

*Choeropsis liberiensis*

- ↔ up to 5 ft (up to 1.5 m)
- 📍 W. Africa
- ⊗ Endangered

The pygmy hippopotamus is much smaller and rarer than its big relative, with a more compact head and narrower feet with fewer webbed toes. A solitary animal, it follows well-worn forest trails to forage at night for leafy foliage and fruit, retreating to a muddy swamp during the day. Like its larger cousin, it has delicate skin that dries out easily in the sun, so it must keep the skin moist by staying close to water.

## Guanaco

*Lama guanicoe*

- ↔ 3–7 ft (0.9–2.1 m)
- 📍 W. to S. South America
- ⊗ Common

This slender, brown-coated relative of the domestic llama is specialized for life in the cold, arid foothills of the Andes, where it is found at altitudes of up to 15,000 ft (4,500 m) or more. It lives in family groups of one breeding male and up to seven adult females and young; unmated males live in separate herds. Guanacos feed mainly on grasses and shrubs, but some also survive in the hostile Atacama Desert, where they eat cacti and lichens.

## Dromedary

*Camelus dromedarius*

- ↔ 7–11 ft (2.1–3.4 m)
- 📍 N. and E. Africa, W. and S. Asia
- ⊗ Not known

Although widespread across north Africa and the Middle East, this one-humped camel is extinct in the wild. The only place where it lives in the wild is Australia, where captive dromedaries, imported to work in the desert, have formed feral breeding populations. The dromedary is superbly adapted for desert life. The hump stores fat that can be broken down into water and energy, allowing the camel to survive for weeks without drinking.

