

NATURAL HABITAT

In its native Amazon, the cane toad is regulated by many predators. As an introduced species in Hawaii and Australia, however, this poisonous amphibian breeds out of control.



CANE TOAD

CLIMBING AROUND

Colorful and pheasant-sized as adult birds, with an impressive headcrest of feathers, as chicks, hoatzins have two claws on each wing, which they use to climb through waterside vegetation until they are able to fly. They lose the claws before adulthood.

HOATZIN CHICK



ENVIRONMENTAL MONITOR

Two species of Amazon river dolphin, the pink river dolphin and the tucuxi, are good indicators of the health of the freshwater ecosystems they inhabit. Fewer dolphins are found where there is existing river degradation, such as poor water quality and overfishing by large human populations.

PINK RIVER DOLPHIN

Contains around 390 billion trees of 16,000 different species

An estimated 2.5 million species of insects live here


LOCATION

The Amazon rainforest stretches across nine South American countries: Brazil, French Guiana, Suriname, Guyana, Venezuela, Colombia, Ecuador, Peru, and Bolivia.



CLIMATE

The Amazon rainforest's climate is tropical and humid. Although all months are wet, rainfall is greatest between December and April, when more than 8 in (200 mm) of rain falls on average each month in Manaus, the region's largest city.



DEFORESTATION

The Amazon rainforest is being felled at such a rate that an estimated 135 plant and animal species become extinct daily. The sustainable harvest of rubber and other wild products, such as Brazil nuts, helps to protect the rainforest from the devastating effects of total land clearance for projects such as soy plantations.

