Table 3.1. STD and their symptoms

Name of the Disease	Causative agent	Symptom	Incubation period
Bacterial STI			
Gonorrhoea	*Neisseria gonorrhoeae*	Affects the urethra, rectum and throat and in females the cervix also get affected. Pain and pus discharge in the genital tract and burning sensation during urination.	2 to 5 days
Syphilis	*Treponema pallidum*	**Primary stage** Formation of painless ulcer on the external genitalia. **Secondary stage** Skin lesions, rashes, swollen joints and fever and hair loss. **Tertiary stage** Appearance of chronic ulcers on nose, lower legs and palate. Loss of movement, mental disorder, visual impairment, heart problems, gummas (soft non-cancerous growths) etc.,	10 to 90 days
Chlamydiasis	*Chlamydia trachomatis*	Trachoma , affects the cells of the columnar epithelium in the urinogenital tract, respiratory tract and conjunctiva.	2 to 3 weeks or upto 6 weeks
Lymphogranuloma venereum	*Chlamydia trachomatis*	Cutaneous or mucosal genital damage, urithritis and endocervicitis. Locally harmful ulcerations and genital elephantiasis.	
Viral STI			
Genital herpes	Herpes simplex virus	Sores in and around the vulva, vagina, urethra in female or sores on or around the penis in male. Pain during urination, bleeding between periods. Swelling in the groin nodes.	2- 21 days (average 6 days)
Genital warts	Human papilloma virus (HPV)	Hard outgrowths (Tumour) on the external genitalia, cervix and perianal region.	1-8 months
Hepatitis-B	Hepatitis B virus (HBV)	Fatigue, jaundice, fever, rashes and stomach pain. Liver cirrhosis and liver failure occur in the later stage.	30-80 days
AIDS	Human immunodeficiency virus (HIV)	Enlarged lymph nodes, prolonged fever, prolonged diarrhoea, weight reduction, night sweating.	2 to 6 weeks even more than 10 years

TN_GOVT_XII_Zoology_chapter 3.indd 37 24-02-2020 4.00.51 PM