Remark

(i) The graph of a periodic function consists of repetitions of the portion of the graph on an interval of length of its period.

(ii) The graph of an odd function is symmetric with respect to the origin and the graph of an even function is symmetric about the y -axis.

4.2.4 Inverse functions

Remember that a function is a rule that, given one value, always gives back a unique value as its answer. For existence, the inverse of a function has to satisfy the above functional requirement. Let us explain this with the help of an example.

Let us consider a set of all human beings not containing identical twins. Every human being from our set, has a blood type and a DNA sequence. These are functions, where a person is the input and the output is blood type or DNA sequence. We know that many people have the same blood type but DNA sequence is unique to each individual. Can we map backwards? For instance, if you know the blood type, do you know specifically which person it came from? The answer is NO. On the other hand, if you know a DNA sequence, a unique individual from our set corresponds to the known DNA sequence. When a function is one-to-one, like the DNA example, then mapping backward is possible. The reverse mapping is called the inverse function. Roughly speaking, the inverse function undoes what the function does.

For any right triangle, given one acute angle and the length of one side, we figure out what the other angles and sides are. But, if we are given only two sides of a right triangle, we need a procedure that leads us from a ratio of sides to an angle. This is where the notion of an inverse to a trigonometric function comes into play.

We know that none of the trigonometric functions is one-to-one over its entire domain. For instance, given $\sin\theta = 0.5$, we have infinitely many $\theta = \frac{\pi}{6}, \frac{5\pi}{6}, \frac{13\pi}{6}, -\frac{7\pi}{6}, -\frac{11\pi}{6}, \cdots$ satisfying the equation. Thus, given $\sin\theta$, it is not possible to recover θ uniquely. To overcome the problem of having multiple angles mapping to the same value, we will restrict our domain suitably before defining the inverse trigonometric function.

To construct the inverse of a trigonometric function, we take an interval small enough such that the function is one-to-one in the restricted interval, but the range of the function restricted to that interval is the whole range. In this chapter, we define the inverses of trigonometric functions with their restricted domains.

4.2.5 Graphs of inverse functions

Assume that f is a bijective function and f^{-1} is the inverse of f. Then, $y = f(x)$ if and only if $x = f^{-1}(y)$. Therefore, (a, b) is a point on the graph of f if and only if (b, a) is the corresponding point on the graph of f^{-1}. This suggests that graph of the inverse function f^{-1} is obtained from the graph of f by interchanging x and y axes. In other words, the graph of f^{-1} is the mirror image of the graph of f in the line $y = x$ or equivalently, the graph of f^{-1} is the reflection of the graph of f in the line $y = x$.

Chapter 4 Inverse Trigonometry.indd 132 31-01-2020 17:40:46