Similarly, $\tan^{-1}\left(\frac{d}{1+a_2a_3}\right) = \tan^{-1}\left(\frac{a_3-a_2}{1+a_2a_3}\right) = \tan^{-1}a_3 - \tan^{-1}a_2$.

Continuing inductively, we get

$\tan^{-1}\left(\frac{d}{1+a_na_{n-1}}\right) = \tan^{-1}\left(\frac{a_n-a_{n-1}}{1+a_{n-1}a_n}\right) = \tan^{-1}a_n - \tan^{-1}a_{n-1}$.

Adding vertically, we get

$\tan^{-1}\left(\frac{d}{1+a_1a_2}\right) + \tan^{-1}\left(\frac{d}{1+a_2a_3}\right) + \ldots + \tan^{-1}\left(\frac{d}{1+a_na_{n-1}}\right) = \tan^{-1}a_n - \tan^{-1}a_1$.

$\tan\left[\tan^{-1}\left(\frac{d}{1+a_1a_2}\right) + \tan^{-1}\left(\frac{d}{1+a_2a_3}\right) + \ldots + \tan^{-1}\left(\frac{d}{1+a_na_{n-1}}\right)\right] = \tan[\tan^{-1}a_n - \tan^{-1}a_1]$

$= \tan\left[\tan^{-1}\left(\frac{a_n-a_1}{1+a_la_n}\right)\right] = \frac{a_n-a_1}{1+a_la_n}$.

Example 4.24 Solve $\tan^{-1}\left(\frac{1-x}{1+x}\right) = \frac{1}{2}\tan^{-1}x$ for $x > 0$.

Solution

$\tan^{-1}\left(\frac{1-x}{1+x}\right) = \frac{1}{2}\tan^{-1}x$ gives $\tan^{-1}1 - \tan^{-1}x = \frac{1}{2}\tan^{-1}x$.

Therefore, $\frac{\pi}{4} = \frac{3}{2}\tan^{-1}x$, which in turn reduces to $\tan^{-1}x = \frac{\pi}{6}$

Thus, $x = \tan\frac{\pi}{6} = \frac{1}{\sqrt{3}}$.

Example 4.25 Solve $\sin^{-1}x > \cos^{-1}x$

Solution

Given that $\sin^{-1}x > \cos^{-1}x$. Note that $-1 \le x \le 1$.

Adding both sides by $\sin^{-1}x$, we get

$\sin^{-1}x + \sin^{-1}x > \cos^{-1}x + \sin^{-1}x$, which reduces to $2\sin^{-1}x > \frac{\pi}{2}$.

As sine function increases in the interval $\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$, we have $x > \sin\frac{\pi}{4}$ or $x > \frac{1}{\sqrt{2}}$.

Thus, the solution set is the interval $\left(\frac{1}{\sqrt{2}}, 1\right]$.

Example 4.26

Show that $\cot(\sin^{-1}x) = \frac{\sqrt{1-x^2}}{x}, -1 \le x \le 1$ and $x \ne 0$

Solution

Let $\sin^{-1}x = \theta$. Then, $x = \sin\theta$ and $x \ne 0$, we get $\theta \in \left[\frac{-\pi}{2}, 0\right) \cup \left(0, \frac{\pi}{2}\right]$.

Hence, $\cos\theta \ge 0$ and $\cos\theta = \sqrt{1-\sin^2\theta} = \sqrt{1-x^2}$.

Thus, $\cot(\sin^{-1}x) = \cot\theta = \frac{\sqrt{1-x^2}}{x}$, $|x| \le 1$ and $x \ne 0$

Chapter 4 Inverse Trigonometry.indd 164 31-01-2020 17:53:07