7. The equation of the normal to the circle $x^2+y^2-2x-2y+1=0$ which is parallel to the line $2x+4y=3$ is

(1) $x+2y=3$ (2) $x+2y+3=0$ (3) $2x+4y+3=0$ (4) $x-2y+3=0$

8. If $P(x,y)$ be any point on $16x^2+25y^2=400$ with foci $F_1(3,0)$ and $F_2(-3,0)$ then PF_1+PF_2 is

(1) 8 (2) 6 (3) 10 (4) 12

9. The radius of the circle passing through the point $(6,2)$ two of whose diameter are $x+y=6$ and $x+2y=4$ is

(1) 10 (2) $2\sqrt{5}$ (3) 6 (4) 4

10. The area of quadrilateral formed with foci of the hyperbolas $\frac{x^2}{a^2}-\frac{y^2}{b^2}=1$ and $\frac{x^2}{a^2}-\frac{y^2}{b^2}=-1$ is

(1) $4(a^2+b^2)$ (2) $2(a^2+b^2)$ (3) a^2+b^2 (4) $\frac{1}{2}(a^2+b^2)$

11. If the normals of the parabola $y^2=4x$ drawn at the end points of its latus rectum are tangents to the circle $(x-3)^2+(y+2)^2=r^2$, then the value of r^2 is

(1) 2 (2) 3 (3) 1 (4) 4

12. If $x+y=k$ is a normal to the parabola $y^2=12x$, then the value of k is

(1) 3 (2) -1 (3) 1 (4) 9

13. The ellipse E_1 : $\frac{x^2}{9}+\frac{y^2}{4}=1$ is inscribed in a rectangle R whose sides are parallel to the coordinate axes. Another ellipse E_2 passing through the point $(0,4)$ circumscribes the rectangle R. The eccentricity of the ellipse is

(1) $\frac{\sqrt{2}}{2}$ (2) $\frac{\sqrt{3}}{2}$ (3) $\frac{1}{2}$ (4) $\frac{3}{4}$

14. Tangents are drawn to the hyperbola $\frac{x^2}{9}-\frac{y^2}{4}=1$ parallel to the straight line $2x-y=1$. One of the points of contact of tangents on the hyperbola is

(1) $\left(\frac{9}{2\sqrt{2}},\frac{-1}{\sqrt{2}}\right)$ (2) $\left(\frac{-9}{2\sqrt{2}},\frac{1}{\sqrt{2}}\right)$ (3) $\left(\frac{9}{2\sqrt{2}},\frac{1}{\sqrt{2}}\right)$ (4) $\left(3\sqrt{3},-2\sqrt{2}\right)$

15. The equation of the circle passing through the foci of the ellipse $\frac{x^2}{16}+\frac{y^2}{9}=1$ having centre at $(0,3)$ is

(1) $x^2+y^2-6y-7=0$ (2) $x^2+y^2-6y+7=0$

(3) $x^2+y^2-6y-5=0$ (4) $x^2+y^2-6y+5=0$

Chapter 5 Analytical Geometry.indd 216 31-01-2020 17:59:29