7. If $\vec{a}$ and $\vec{b}$ are unit vectors such that $[\vec{a}, \vec{b}, \vec{a}\times\vec{b}]=\frac{1}{4}$, then the angle between $\vec{a}$ and $\vec{b}$ is

(1) $\frac{\pi}{6}$ (2) $\frac{\pi}{4}$ (3) $\frac{\pi}{3}$ (4) $\frac{\pi}{2}$

8. If $\vec{a}=\hat{i}+\hat{j}+\hat{k}$, $\vec{b}=\hat{i}+\hat{j}$, $\vec{c}=\hat{i}$ and $(\vec{a}\times\vec{b})\times\vec{c}=\lambda\vec{a}+\mu\vec{b}$, then the value of $\lambda+\mu$ is

(1) 0 (2) 1 (3) 6 (4) 3

9. If $\vec{a},\vec{b},\vec{c}$ are non-coplanar, non-zero vectors such that $[\vec{a}, \vec{b}, \vec{c}]=3$, then $\{[\vec{a}\times\vec{b}, \vec{b}\times\vec{c}, \vec{c}\times\vec{a}]\}^2$ is equal to

(1) 81 (2) 9 (3) 27 (4)18

10. If $\vec{a},\vec{b},\vec{c}$ are three non-coplanar unit vectors such that $\vec{a}\times(\vec{b}\times\vec{c})=\frac{\vec{b}+\vec{c}}{\sqrt{2}}$, then the angle between $\vec{a}$ and $\vec{b}$ is

(1) $\frac{\pi}{2}$ (2) $\frac{3\pi}{4}$ (3) $\frac{\pi}{4}$ (4) π

11. If the volume of the parallelepiped with $\vec{a}\times\vec{b}$, $\vec{b}\times\vec{c}$, $\vec{c}\times\vec{a}$ as coterminous edges is 8 cubic units, then the volume of the parallelepiped with $(\vec{a}\times\vec{b})\times(\vec{b}\times\vec{c}),(\vec{b}\times\vec{c})\times(\vec{c}\times\vec{a})$ and $(\vec{c}\times\vec{a})\times(\vec{a}\times\vec{b})$ as coterminous edges is,

(1) 8 cubic units (2) 512 cubic units (3) 64 cubic units (4) 24 cubic units

12. Consider the vectors $\vec{a},\vec{b},\vec{c},\vec{d}$ such that $(\vec{a}\times\vec{b})\times(\vec{c}\times\vec{d})=\vec{0}$. Let P_1 and P_2 be the planes determined by the pairs of vectors $\vec{a},\vec{b}$ and $\vec{c},\vec{d}$ respectively. Then the angle between P_1 and P_2 is

(1) 0° (2) 45° (3) 60° (4) 90°

13. If $\vec{a}\times(\vec{b}\times\vec{c})=(\vec{a}\times\vec{b})\times\vec{c}$, where $\vec{a},\vec{b},\vec{c}$ are any three vectors such that $\vec{b}\cdot\vec{c}\neq 0$ and $\vec{a}\cdot\vec{b}\neq 0$, then $\vec{a}$ and $\vec{c}$ are

(1) perpendicular (2) parallel

(3) inclined at an angle $\frac{\pi}{3}$ (4) inclined at an angle $\frac{\pi}{6}$

14. If $\vec{a}=2\hat{i}+3\hat{j}-\hat{k}$, $\vec{b}=\hat{i}+2\hat{j}-5\hat{k}$, $\vec{c}=3\hat{i}+5\hat{j}-\hat{k}$, then a vector perpendicular to $\vec{a}$ and lies in the plane containing $\vec{b}$ and $\vec{c}$ is

(1) $-17\hat{i}+21\hat{j}-97\hat{k}$ (2) $17\hat{i}+21\hat{j}-123\hat{k}$

(3) $-17\hat{i}-21\hat{j}+97\hat{k}$ (4) $-17\hat{i}-21\hat{j}-97\hat{k}$

15. The angle between the lines $\frac{x-2}{3}=\frac{y+1}{-2}, z=2$ and $\frac{x-1}{1}=\frac{2y+3}{3}=\frac{z+5}{2}$ is

(1) $\frac{\pi}{6}$ (2) $\frac{\pi}{4}$ (3) $\frac{\pi}{3}$ (4) $\frac{\pi}{2}$

Chapter 6 Vector Algebra.indd 277 31-01-2020 17:53:55