20. Which of the following is/are correct?

(i) Adjoint of a symmetric matrix is also a symmetric matrix.

(ii) Adjoint of a diagonal matrix is also a diagonal matrix.

(iii) If A is a square matrix of order n and λ is a scalar, then $\text{adj}(\lambda A) = \lambda^n \text{adj}(A)$.

(iv) $A(\text{adj}A) = (\text{adj}A)A = |A|I$

(1) Only (i) (2) (ii) and (iii) (3) (iii) and (iv) (4) (i), (ii) and (iv)

21. If $\rho(A) = \rho([A|B])$, then the system $AX = B$ of linear equations is

(1) consistent and has a unique solution (2) consistent

(3) consistent and has infinitely many solution (4) inconsistent

22. If $0 \le \theta \le \pi$ and the system of equations $x + (\sin\theta)y - (\cos\theta)z = 0, (\cos\theta)x - y + z = 0,$ $(\sin\theta)x + y - z = 0$ has a non-trivial solution then θ is

(1) $\frac{2\pi}{3}$ (2) $\frac{3\pi}{4}$ (3) $\frac{5\pi}{6}$ (4) $\frac{\pi}{4}$

23. The augmented matrix of a system of linear equations is $\begin{bmatrix} 1 & 2 & 7 & 3 \\ 0 & 1 & 4 & 6 \\ 0 & 0 & \lambda-7 & \mu+5 \end{bmatrix}$. The system has infinitely many solutions if

(1) $\lambda = 7, \mu \neq -5$ (2) $\lambda = -7, \mu = 5$ (3) $\lambda \neq 7, \mu \neq -5$ (4) $\lambda = 7, \mu = -5$

24. Let $A = \begin{bmatrix} 2 & -1 & 1 \\ -1 & 2 & -1 \\ 1 & -1 & 2 \end{bmatrix}$ and $4B = \begin{bmatrix} 3 & 1 & -1 \\ 1 & 3 & x \\ -1 & 1 & 3 \end{bmatrix}$. If B is the inverse of A, then the value of x is

(1) 2 (2) 4 (3) 3 (4) 1

25. If $A = \begin{bmatrix} 3 & -3 & 4 \\ 2 & -3 & 4 \\ 0 & -1 & 1 \end{bmatrix}$, then $\text{adj}(\text{adj}\, A)$ is

(1) $\begin{bmatrix} 3 & -3 & 4 \\ 2 & -3 & 4 \\ 0 & -1 & 1 \end{bmatrix}$ (2) $\begin{bmatrix} 6 & -6 & 8 \\ 4 & -6 & 8 \\ 0 & -2 & 2 \end{bmatrix}$ (3) $\begin{bmatrix} -3 & 3 & -4 \\ -2 & 3 & -4 \\ 0 & 1 & -1 \end{bmatrix}$ (4) $\begin{bmatrix} 3 & -3 & 4 \\ 0 & -1 & 1 \\ 2 & -3 & 4 \end{bmatrix}$

SUMMARY

(1) Adjoint of a square matrix A = Transpose of the cofactor matrix of A.

(2) $A(\text{adj}\, A) = (\text{adj}\, A)A = |A| I_n$.

(3) $A^{-1} = \frac{1}{|A|} \text{adj}\, A$.

(4) (i) $|A^{-1}| = \frac{1}{|A|}$ (ii) $(A^T)^{-1} = (A^{-1})^T$ (iii) $(\lambda A)^{-1} = \frac{1}{\lambda} A^{-1}$, where λ is a non-zero scalar.

Chapter 1 Matrices.indd 50 31-01-2020 17:17:12