II-VOLUMETRIC ANALYSIS

1. Estimation of Ferrous Sulphate (Fe^{2+})

Aim :

To estimate the amount of ferrous sulphate dissolved in 750 ml of the given unknown solution volumetrically. For this you are given with a standard solution of ferrous ammonium sulphate (FAS) of normality 0.1102 N and potassium permanganate solution as link solution.

Principle:

During these titrations, Fe^{2+} ions (from ferrous salts) are oxidised to MnO_4^- ions and MnO_4^- ion (from Mn^{2+}) is reduced to Mn^{2+} ion.

Oxidation : $5\ Fe^{2+} \longrightarrow 5\ Fe^{3+} + 5e^-$

Reduction : $\underset{\text{Pink}}{MnO_4^-} + 8H^+ + 5e^- \longrightarrow \underset{\text{colourless}}{Mn^{2+}} + 4H_2O$

Overall reaction : $5Fe^{2+} + MnO_4^- + 8H^+ \longrightarrow 5Fe^{3+} + Mn^{2+} + 4H_2O$

Short procedure:

S.no	Content	Titration-I	Titration-II
1	Burette solution	$KMnO_4$	$KMnO_4$
2	Pipette solution	20 ml of standard FAS	20 ml of unknown $FeSO_4$
3	Acid to be added	20ml of 2N H_2SO_4 (approx)	20ml of 2N H_2SO_4 (approx)
4	Temperature	Lab temperature	Lab temperature
5	Indicator	Self-indicator ($KMnO_4$)	Self-indicator ($KMnO_4$)
6	End point	Appearance of permanent pale pink colour	Appearance of permanent pale pink colour
7	Equivalent weight of $FeSO_4$ = 278		

Procedure :

Titration–I

(Link $KMnO_4$)Vs (Standard FAS)

Burette is washed with water, rinsed with $KMnO_4$ solution and filled with same $KMnO_4$ solution up to the zero mark. Exactly 20 ml of standard FAS solution is pipetted out into the clean, washed conical flask. To this FAS solution, approximately 20ml of 2N sulphuric acid is added. This mixture is titrated against $KMnO_4$ Link solution from the burette. $KMnO_4$ is added drop wise till the appearance of permanent pale pink colour. Burette reading is noted, and the same procedure is repeated to get concordant values.

organic analysis.indd 265 19-01-2021 12:24:22