$$r_2 = r\left(1+\frac{2a\cos\theta}{r}\right)^{\frac{1}{2}}$$

Using Binomial theorem, we get

$$\frac{1}{r_2} = \frac{1}{r}\left(1-a\frac{\cos\theta}{r}\right) \qquad (1.37)$$

Substituting equation (1.37) and (1.36) in equation (1.35),

$$V = \frac{1}{4\pi\epsilon_\circ} q\left(\frac{1}{r}\left(1+a\frac{\cos\theta}{r}\right) - \frac{1}{r}\left(1-a\frac{\cos\theta}{r}\right)\right)$$

$$V = \frac{q}{4\pi\epsilon_\circ}\left(\frac{1}{r}\left(1+a\frac{\cos\theta}{r}-1+a\frac{\cos\theta}{r}\right)\right)$$

$$V = \frac{1}{4\pi\epsilon_\circ}\frac{2aq}{r^2}\cos\theta$$

But the electric dipole moment $p = 2qa$ and we get,

$$V = \frac{1}{4\pi\epsilon_0}\left(\frac{p\cos\theta}{r^2}\right)$$

Now we can write $p\cos\theta = \vec{p}\cdot\hat{r}$, where $\hat{r}$ is the unit vector from the point O to point P. Hence the electric potential at a point P due to an electric dipole is given by

$$V = \frac{1}{4\pi\epsilon_\circ}\frac{\vec{p}\cdot\hat{r}}{r^2} \quad (r>>a) \qquad (1.38)$$

Equation (1.38) is valid for distances very large compared to the size of the dipole. But for a point dipole, the equation (1.38) is valid for any distance.

Special cases

Case (i) If the point P lies on the axial line of the dipole on the side of $+q$, then $\theta = 0$. Then the electric potential becomes

$$V = \frac{1}{4\pi\epsilon_\circ}\frac{p}{r^2} \qquad (1.39)$$

Case (ii) If the point P lies on the axial line of the dipole on the side of $-q$, then $\theta = 180^\circ$. Then

$$V = -\frac{1}{4\pi\epsilon_\circ}\frac{p}{r^2} \qquad (1.40)$$

Case (iii) If the point P lies on the equatorial line of the dipole, then $\theta = 90^\circ$. Hence

$$V = 0 \qquad (1.41)$$

Note The potential due to an electric dipole falls as $\frac{1}{r^2}$ and the potential due to a single point charge falls as $\frac{1}{r}$. Thus the potential due to the dipole falls faster than that due to a monopole (point charge). As the distance increases from electric dipole, the effects of positive and negative charges nullify each other

1.5.4 Equi-potential Surface

Consider a point charge q located at some point in space and an imaginary sphere of radius r is chosen by keeping the charge q at its centre (Figure 1.26(a)). The electric potential at all points on the surface of the given sphere is the same. Such a surface is called an equipotential surface.

An equipotential surface is a surface on which all the points are at the same electric potential. For a point charge the equipotential surfaces are concentric spherical surfaces as shown in Figure 1.26(b). Each spherical surface is an equipotential surface but the

UNIT-1(XII-Physics_Vol-1).indd 31 14-02-2020 19:32:57