
**(iii)** To arrive at equation (1.62), we have chosen a spherical surface. This imaginary surface is called a Gaussian surface. The shape of the Gaussian surface to be chosen depends on the type of charge configuration and the kind of symmetry existing in that charge configuration. The electric field is spherically symmetric for a point charge, therefore spherical Gaussian surface is chosen. Cylindrical and planar Gaussian surfaces can be chosen for other kinds of charge configurations.

**(iv)** In the LHS of equation (1.62), the electric field $\vec{E}$ is due to charges present inside and outside the Gaussian surface but the charge $Q_{encl}$ denotes the charges which lie only inside the Gaussian surface.

## EXAMPLE 1.18




**(i)** In figure (a), calculate the electric flux through the closed areas $A_1$ and $A_2$.

**(ii)** In figure (b), calculate the electric flux through the cube

### *Solution*

**(i)** In figure (a), the area $A_1$ encloses the charge $Q$. So electric flux through this closed surface $A_1$ is $\frac{Q}{\epsilon_0}$. But the closed surface $A_2$ contains no charges inside, so electric flux through $A_2$ is zero.

**(ii)** In figure (b), the net charge inside the cube is $3q$ and the total electric flux in the cube is therefore $\Phi_E = \frac{3q}{\epsilon_0}$. Note that the charge -10 $q$ lies outside the cube and it will not contribute the total flux through the surface of the cube.

## 1.6.4 Applications of Gauss law

Electric field due to any arbitrary charge configuration can be calculated using Coulomb's law or Gauss law. If the charge configuration possesses some kind of symmetry, then Gauss law is a very efficient way to calculate the electric field. It is illustrated in the following cases.

**(i) Electric field due to an infinitely long charged wire**

Consider an infinitely long straight wire having uniform linear charge density $\lambda$(charge per unit length). Let P be a point located at a perpendicular distance $r$ from the wire (Figure 1.36(a)). The electric field at the point P can be found using Gauss law.

We choose two small charge elements $A_1$ and $A_2$ on the wire which are at equal distances from the point P. The resultant electric field due to these two charge elements points radially away from the charged wire and the magnitude of electric field is same at all points on the circle of radius r. This is shown in the Figure 1.36(b). Since the charged wire possesses



UNIT-1(XII-Physics_Vol-1).indd 41 14-02-2020 19:33:17