
AIDS	Human Immuno deficiency Virus (HIV)	Direct contact with infected blood, semen or vaginal fluids	12 years	flu-like symptoms such as fever, sore throat and fatigue after few weeks of infection	anti-retroviral treatment (ARVT)

10.2.3. Fungal Infections

Fungi are eukaryotic protista, recognized as causative agents of human disease earlier than bacteria. Fungal infections (mycosis) are most common among those patients who use antibiotics for prolonged period of time. These antibiotics not only kill pathogenic bacteria but also target the normal flora of human body (useful bacteria) and give rise to fungal growth. Human fungal infections are usually of two types: superficial and deep infection. Fungi causing superficial mycoses are specialized saprophytes, with the capacity to digest keratin. Superficial mycoses are of two types - surface infections (only on dead layers of skin) and cutaneous infections (cornified layer).



Mucormycosis molds



Histoplasma fungi

Disease Name	Pathogen	Mode of transmission	Symptoms	Therapy
Candidiasis or yeast infections	*Candida* yeasts *Candida albicans*	Direct contact	itching and swelling, redness and soreness	Nystatin , Clotrimazole and fluconazole
Jock itch or Tinea cruris (Dermatophytoses)	*Trichophyton rubrum* and *T. mentagrophytes*	direct contact with an infected person	redness in the groin, buttocks, or thighs, itching	miconazole, clotrimazole, ketoconazole

XII U10 Immunology.indd 235 4/8/2020 3:27:55 PM