3.	Heterophile or Heterogenic or Cross reactive antigen	These antigens interact with antibodies produced against another antigen.	Forssman antigen. These antigens are present on mucosal cells of GI tract of human and RBCs of horse.
4.	Mitogenic antigens	Since some mitogens (they induce cell division) activate immune cells, they can be considered as mitogenic antigens.	Lipopolysaccharides induce B cell proliferation, Concanavalin – A (CON A) and Phytohemagglutinin (PHA) induce T cell proliferation and Pokeweed mitogen (PWM) induces both T and B cell proliferation.
5.	Superantigens	These antigens activate T cells non-specifically.	TSST-1 (Toxic shock syndrome toxin – 1) produced from *Staphylococcus sp*. . - Exogenous superantigens Mouse mammary tumor viral (MMTV) antigens - Endogenous superantigens
6.	Exogenous antigens	These antigens are usually secretory products of microbes or soluble antigens. They are usually present outside the host cell	Diphtheria toxin from *Cornebacterium diptheriae*
7.	Endogenous antigens	They are usually microbes which are present inside the cell or particulate antigen.	Virus and endotoxins

Sl. No.	**Types of Antigens**	**Nature**	**Example**
8.	T cell dependent antigens	These antigens require T cells especially T_H cells to induce immune response.	Soluble antigens

XII U10 Immunology.indd 245 4/8/2020 3:27:59 PM