Titration: 1 Sodium hydroxide vs oxalic acid

S.No.	volume of solution in conical flask	Burette readings (ml)		Volume of NaOH consumed (ml)	Concordant volume
		Initial	Final		
	10	0			
	10	0			

Titration : 2 Blank titration (sodium hydroxide vs formaldehyde)

S.No.	volume of solution in conical flask	Burette readings (ml)		Volume of NaOH consumed (ml)	Concordant volume
		Initial	Final		
	20	0			
	20	0			

Titration : 3 Sodium hydroxide vs unknow solution

S.No.	volume of solution in conical flask	Burette readings (ml)		Volume of NaOH consumed (ml)	Concordant volume
		Initial	Final		
	30	0			
	30	0			

Calculation

Titration 1

Volume of oxalic acid (V1) = 10ml
Normality of oxalic acid (N1) = 0.1N
Volume of NaOH (V2) =
Normality of NaOH (N2) = N1V1 / V2

Titration 2

Volume of NaOH (V3) =

Titration 3

Volume of NaOH (V4) =
Net Volume of NaOH (V5) = V4 – V3

Normality of NaOH (N2) =

Volume of amino acid (glycine) (V6) = 10ml
Normality of amino acid (glycine) = N2 V5 / V6
Normality = (Weight / L) / equivalent weight
Weight /L = Normality x equivalent weight
= Z gram of glycine/1000ml (1L)
Therefore, 100ml of the test solution (glycine) will contain 100 x $\frac{Z}{1000}$ gram of glycine

Result:

The amount of glycine present in 100ml of the given solution was found to be ____ g.

Practical_Bio-Chem-EM.indd 288 4/8/2020 3:28:53 PM