- **Polyphagia**: Excessive appetite leads to polyphagia and increased intake of food. This is to replace the lost nourishment. The diabetic has voracious appetite, but inspite of over eating, they lose weight and become lean and emaciated.
- Presence of ketones in the urine (ketones are a byproduct of the breakdown of muscle and fat that happens when there's not enough available insulin)
- Fatigue
- Irritability
- Blurred vision
- Slow-healing sores
- Frequent yeast or urinary tract infections

3.8.3. Diagnosis

Diabetes is diagnosed with fasting sugar blood tests or with A1C blood tests, also known as **glycated hemoglobin tests**. You do not have to be in fasting for an A1C blood test. A fasting blood sugar test is performed after you have had nothing to eat or drink for at least eight hours. Normal fasting blood sugar is less than **100 mg/dl (5.6 mmol/l)**.

Test	Normal glucose level	Level of glucose in diabetes patient
Fasting Glucose Test	Less than 100 mg/dl	126 mg/dl or higher
Random (anytime) Glucose Test	Less than 140 mg/dl	200 mg/dl or higher
A1C Test	Less than 5.7%	6.5% or higher

3.8.3.1. Glucose tolerance test

The glucose tolerance test, also known as the **oral glucose tolerance test** (OGTT), measures your body's response to sugar (glucose). The glucose tolerance test can be used to screen for type 2 diabetes.

Before the test begins, a sample of blood will be taken.

XII U3 Carbohydrates.indd 86 4/8/2020 3:06:55 PM