


Fig. 1.5 Strobilation in *Aurelia*

(**Fig. 1.5**). In the process of strobilation, several transverse fissions occur simultaneously giving rise to a number of individuals which often do not separate immediately from each other e.g. *Aurelia*. **Plasmotomy** is the division of multinucleated parent into many multinucleate daughter individuals with the division of nuclei. Nuclear division occurs later to maintain normal number of nuclei. Plasmotomy occurs in *Opalina* and *Pelomyxa (*Giant *Amoeba).*

During unfavourable conditions *Amoeba* multiplies by **sporulation** without encystment. Nucleus breaks into several small fragments or chromatin blocks. Each fragment develops a nuclear membrane, becomes surrounded by cytoplasm and develops a spore-case around it (**Fig. 1.6**). When conditions become favourable, the parent body disintegrates and the spores are liberated, each hatching into a young amoeba.

In **budding**, the parent body produces one or more buds and each bud grows into a young one. The buds separate from the parent to lead a normal life. In sponges, the buds constrict and detach from the parent body and the bud develops into a new sponge (**Fig. 1.7**).




Fig. 1.6 Sporulation in *Amoeba*

TN_GOVT_XII_Bio_Zoology_EM_chapter 1.indd 4 20-03-2020 10.31.09 AM