**Table 7.7 The cellular composition of adult human blood**

| Cell type | Number of cells per µl | Approximate percentage |
|---|---|---|
| Red blood cells | 4200,000 - 6500,000 | - |
| **White blood cells** | | |
| Agranulocytes | | |
| Lymphocytes | 1500 - 4000 | 20-30 |
| Monocytes | 200 - 950 | 2-7 |
| Granulocytes | | |
| Neutrophils | 2000-7000 | 50-70 |
| Basophils | 50-100 | <1 |
| Eosinophils | 40-500 | 2-5 |
| Platelets | 150,000-500,000 | - |

**Lymphocytes**

About 20-30% of the white blood cells are lymphocytes. They have a large nucleus filling most of the cell, surrounded by a little cytoplasm. The two main types of lymphocytes are B and T lymphocytes. Both these are produced in the bone marrow. B lymphocytes (B cells) stay in the bone marrow until they are mature. Then they circulate around the body. Some remain in the blood, while others accumulate in the lymph nodes and spleen. T lymphocytes leave the bone marrow and mature in the thymus gland. Once mature, T cells also accumulate in the same areas of the body as B cells. Lymphocytes have receptor proteins on their surface. When receptors on a B cell bind with an antigen, the B cell becomes activated and divides rapidly to produce plasma cells. The plasma cells produce antibodies. Some B cells do not produce antibodies but become memory cells. These cells are responsible for secondary immune response. T lymphocytes do not produce antibodies. They recognize antigen-presenting cells and destroy them. The two important types of T cells are Helper T cells and Killer T cells. Helper T cells release a chemical





**Fig. 7.14 Cells of the immune system**



TN_GOVT_XII_Bio_Zoology_EM_chapter 7.indd 122 20-03-2020 10.34.19 AM