Somatic cell therapy involves the insertion of a fully functional and expressible gene into a target somatic cell to correct a genetic disease permanently whereas Germline gene therapy involves the introduction of DNA into germ cells which is passed on to the successive generations. Gene therapy involves isolation of a specific gene and making its copies and inserting them into target cells to make the desired proteins. It is absolutely essential for gene therapists to ensure that the gene is harmless to the patient and it is appropriately expressed and that the body's immune system does not react to the foreign proteins produced by the new genes.

9.3 Stem Cell Therapy

Stem cells are undifferentiated cells found in most of the multi cellular animals. These cells maintain their undifferentiated state even after undergoing numerous mitotic divisions.

Stem cell research has the potential to revolutionize the future of medicine with the ability to regenerate damaged and diseased organs. Stem cells are capable of self renewal and exhibit 'cellular potency'. Stem cells can differentiate into all types of cells that are derived from any of the three germ layers ectoderm, endoderm and mesoderm.

Table 9.1 differentiation between somatic cell gene therapy and germ line gene therapy

SOMATIC CELL GENE THERAPY	GERM LINE GENE THERAPY
Therapeutic genes transferred into the somatic cells.	Therapeutic genes transferred into the germ cells.
Introduction of genes into bone marrow cells, blood cells, skin cells etc.,	Genes introduced into eggs and sperms.
Will not be inherited in later generations.	Heritable and passed on to later generations.

The first clinical gene therapy was given in 1990 by French Anderson to a four year old girl with adenosine deaminase (ADA) deficiency. ADA deficiency or SCID (Severe combined immunodeficiency) is an autosomal recessive metabolic disorder. It is caused by the deletion or dysfunction of the gene coding for ADA enzyme. In these patients the nonfunctioning T-Lymphocytes cannot elicit immune responses against invading pathogens. The right approach for SCID treatment would be to give the patient a functioning ADA which breaks down toxic biological products.

In some children ADA deficiency could be cured by bone marrow transplantation, where defective immune cells could be replaced with healthy immune cells from a donor. In some patients it can be treated by enzyme replacement therapy, in which functional ADA is injected into the patient.

During gene therapy the lymphocytes from the blood of the patient are removed and grown in a nutrient culture medium. A healthy and functional human gene, ADA cDNA encoding this enzyme is introduced into the lymphocytes using a retrovirus. The genetically engineered lymphocytes are subsequently returned to the patient. Since these cells are not immortal, the patient requires periodic infusion of such genetically engineered lymphocytes. The disease could be cured permanently if the gene for ADA isolated from bone marrow cells are introduced into the cells of the early embryonic stages.

TN_GOVT_XII_Bio_Zoology_EM_chapter 9.indd 151 20-03-2020 10.36.15 AM