is the ectoderm. The cells remaining in between the epiblast and the endoderm form the mesoderm. Thus the transformation of the blastocyst into a gastrula with the primary germ layers by the movement of the blastomeres is called **gastrulation**. Each germ layer gives rise to specific tissues, organs and organ systems during organogenesis.

The extra embryonic membranes namely the **amnion, yolk sac, allantois** and **chorion** protect the embryo from dessication, mechanical shock and help in the absorption of nutrients and exchange of gases **(Fig. 2.12)**. The amnion is a double layered translucent membrane filled with the amniotic fluid. It provides a buoyant environment to protect the developing embryo from injury, regulates the temperature of the foetus and provides a medium in which the foetus can move. The yolk sac forms a part of the gut and is the source of the earliest blood cells and blood vessels.




Fig. 2.12 Human foetus within the uterus

The allantois forms a small out pocketing of embryonic tissue at the caudal end of the yolk sac. It is the structural base for the umbilical cord that links the embryo to the placenta and ultimately it becomes part of the urinary bladder. The chorion is the outermost membrane which encloses the embryo and all other membranes and also helps in the formation of the placenta.

The trophoblast cells in the blastocyst send out several finger like projections called **chorionic villi** carrying foetal blood and are surrounded by sinuses that contain maternal blood. The chorionic villi and the uterine tissues form the disc-shaped placenta. **Placenta** is a temporary endocrine organ formed during pregnancy and it connects the foetus to the uterine wall through the umbilical cord. It is the organ by which the nutritive, respiratory and excretory functions are fulfilled. The embryo's heart develops during the fourth week of pregnancy and circulates blood through the umbilical cord and placenta as well as through its own tissues.

The primary germ layers serve as the primitive tissues from which all body organs develop. The ectoderm gives rise to the central nervous system (brain and spinal cord), peripheral nervous system, epidermis and its derivatives and mammary glands. The connective tissue, cartilage and bone, muscles, organs of urinogenital system (kidney, ureter and gonads) arise from the mesoderm. The endodermal derivatives are **e**pithelium of gastrointestinal and respiratory tract, liver, pancreas, thyroid and parathyroids.

Human pregnancy lasts for about 280 days or 40 weeks and is called the **gestation period**. It can be divided for convenience into three trimesters of three months each. The **first trimester** is the main period of organogenesis, the body organs namely the heart, limbs, lungs, liver and external genital organs are well developed. By the end of the **second trimester**, the face is well formed with features, eyelids and eyelashes, eyes blink, body is covered with fine hair, muscle tissue develops and bones become harder. The foetus is fully developed and is ready for delivery by the end of nine months (**third trimester**).

During pregnancy, the placenta acts as a temporary endocrine gland and produces large quantities of **human Chorionic Gonadotropin (hCG), human Chorionic Somatomammotropin (hCS) or human**

TN_GOVT_XII_Bio_Zoology_EM_chapter 2.indd 23 20-03-2020 10.32.03 AM