on the recommendations of **Justice Verma Committee**, 2013 aims at creating a safe and secure environment for both females and males.

3.4. Population explosion and birth control

Increased health facilities and better living conditions have enhanced longevity. According to a recent report from the UN, India's population has already reached 1.26 billion and is expected to become the largest country in population size, surpassing China around 2022. To overcome the problem of population explosion, birth control is the only available solution. People should be motivated to have smaller families by using various contraceptive devices. Advertisements by the Government in the media as well as posters/bills, etc., with a slogan *Naam iruvar namakku iruvar* (we two, ours two) and *Naam iruvar namakku oruvar* (we two, ours one) have also motivated to control population growth in Tamilnadu. Statutory rising of marriageable age of the female to 18 years and that of males to 21 years and incentives given to couples with small families are the other measures taken to control population growth in our country.

Birth control methods

The voluntary use of contraceptive procedures to prevent fertilization or prevent implantation of a fertilized egg in the uterus is termed as birth control. **An ideal contraceptive should be** user friendly, easily available, with least side effects and should not interfere with sexual drive. The contraceptive methods are of two types – **temporary** and **permanent.** Natural, chemical, mechanical and hormonal barrier methods are the temporary birth control methods.

1. Natural method is used to prevent meeting of sperm with ovum. i.e., Rhythm method (safe period), coitus interruptus, continuous abstinence and lactational amenorrhoea.

a. Periodic abstinence/rhythm method Ovulation occurs at about the 14th day of the menstrual cycle. Ovum survives for about two days and sperm remains alive for about 72 hours in the female reproductive tract. Coitus is to be avoided during this time.

b. Continuous abstinence is the simplest and most reliable way to avoid pregnancy is not to have coitus for a defined period that facilitates conception.

c. Coitus interruptus is the oldest family planning method. The male partner withdraws his penis before ejaculation, thereby preventing deposition of semen into the vagina.

d. Lactational amenorrhoea Menstrual cycles resume as early as 6 to 8 weeks from parturition. However, the reappearance of normal ovarian cycles may be delayed for six months during breast-feeding. This delay in ovarian cycles is called **lactational amenorrhoea.** It serves as a natural, but an unreliable form of birth control. Suckling by the baby during breast-feeding stimulates the pituitary to secrete increased prolactin hormone in order to increase milk production. This high prolactin concentration in the mother's blood may prevent menstrual cycle by suppressing the release of GnRH (Gonadotropin Releasing Hormone) from hypothalamus and gonadotropin secretion from the pituitary.

2. Barrier methods In these methods, the ovum and sperm are prevented from meeting so that fertilization does not occur.

a. Chemical barrier Foaming tablets, melting suppositories, jellies and creams are used as chemical agents that inactivate the sperms in the vagina.

b. Mechanical barrier Condoms are a thin sheath used to cover the penis in

TN_GOVT_XII_Bio_Zoology_EM_chapter 3.indd 34 20-03-2020 10.32.44 AM