


Figure 8.13: Urine in Black water Malaria

4. Anaemia: An individual suffering from an attack of malaria, after a few paroxysms becomes temporarily anaemic. The reduction in red blood cells is greater in *P. falciparum* infection than in infection with *P. vivax* and *P. malariae*. This is because *P. falciparum* invades young and mature erythrocytes and the infection rate of red blood cells is also greater.

HOTS

Which stage is infective in Blood transfusion malaria?

8.5.10 Recrudescence

In *P. falciparum* and *P. malariae* infections after the primary attack, sometimes there is a period of latency, during which there is no clinical illness. But some parasites persist in some erythrocytes and gradually increase in numbers. Fresh malarial attacks then develop. It appears after a period of latency usually within weeks after the primary attacks. Persistence of the erythrocytic cycle of the parasites are called recrudescences. In *P. falciparum* infections, recrudescences are seen for 1–2 years, while in *P. malariae* infection, they may last for long periods, even upto 50 years.

Infobits

The global technical strategy for malaria 2016-2030 was adopted by the World health Assembly in May 2015. It provides a comprehensive framework to guide countries in their efforts to accelerate progress towards malaria elimination. The strategy sets the target of reducing global malaria incidence and mortality rates by at least 90% by 2030.

8.5.11 Plasmodium vivax

P. vivax shows a similar life cycle in humans and mosquitoes like that of *P. falciparum*. Except in *P. vivax*, a latent tissue stage, the hypnozoites present in the liver parenchyma.

Relapse in vivax malaria is caused by these hypnozoites. Hypnozoites are the dormant stages of the parasites. These are single – nucleated parasites measuring 4µm–6µm in diameter. These become active and develop into tissue schizonts after a short period of dormancy. This relapse may occur at intervals up to 3 years or more after the first attack. *P. vivax* merozoites invade only young erythrocytes and the reticulocytes.

8.5.12 Clinical Manifestations

P. vivax is the most wide spread species causing malaria in man. However, unlike

TN_GOVT_XII_Micro_Biology_CH08.indd 177 27-02-2019 14:53:05