Serology

Antibodies develop within a few days of infection and **rise in titre of antibodies** may be demonstrated by ELISA, neutralization or complement fixation tests.

Chemotherapy

Indoxyuridine used topically in eye and skin infection, acyclovir and vidarabine are given for deep and systemic infections.

2. Varicella Zoster

In 1889, Von Bokay had suggested that varicella (Chicken pox) and herpes zoster are **different manifestations** of the same virus infection. The virus is therefore called **Varicella zoster virus** (VZV). The chicken pox follows **primary infection** in a non immune individual, while herpes zoster is a **reaction of the latent virus** when the immunity has fallen to infective levels.

VZV is **similar** to the herpes simplex virus in its morphology. It can be grown in cultures of **human fibroblasts human amnion** or HeLa cells. Chicken pox is one of the mildest and **most common of child hood infections**. The disease may, occur at any age.



Herpes gladiatorum is **spread** through skin-on-skin contact. If you kiss someone with a herpes cold sore on their lips, you could become infected.

3. Cytomegaloviruses

Cytomegaloviruses (CMV) formerly known as **salivary gland viruses** are a group of ubiquitous herpes viruses of humans and animals. They are characterized by **enlargement of infected cells** and intranuclear inclusions. In 1926, cytomegalia presumed to be due to viral infection was reported in the salivary glands of guinea pigs and children and the viral agent was called the **'salivary gland virus'.**

Infobits

CMV is the **largest viruses** in the herpes virus family, being 150–200 nm in size.

4. Epstein – Barr Virus

A number of different viruses apparently **'Passenger Viruses'** were isolated from cultured lymphoma cells. **Epstein, Barr and Achong in 1964** observed a new type of herpes virus named it has **'EB Virus'** affecting cells of B lymphocyte. Only human and some sub human primate **B cells have receptors** (CD21 molecules) for the virus.

The source of infection is usually the **saliva of infected persons** who shed the virus in orophryngeal secretions. Intimate oral contact, as in kissing, appears to be the predominant mode of transmission. This accounts for infectious mononucleosis being called as **'The kissing disease'**.

5. Human Herpes Virus Types 6,7,8

A herpes virus, first isolated in 1986 from the **peripheral blood** of patients with lympho proliforative disease called as **human B lymph tropic virus**, renamed as HHV - 6. HHV- 7 was isolated in 1990 from **peripheral CD4 cells** of a healthy

TN_GOVT_XII_Micro_Biology_CH10.indd 214 04-03-2019 12:27:07