


Figure 11.10: Type IV delayed hypersensitivity

and macrophages and cause the bulk damage (Figure 11.10).

Tuberculin reaction (Mantoux Reaction)

When a small dose of tuberculin is injected intra dermally in an individual already having tubercle bacilli, the reaction occurs. It is due to the interaction of sensitized T cell and tubercle bacterium. The reaction is manifested on the skin very late only after 48–72 hours.

11.4 Transplantation

Transfer of living cells, tissues or organs from one part of the body to another or from one individual to another is known as transplantation.

A tissue or organ that is removed from one site and placed to another site usually in a same or different individual is called graft. The individual who provides the graft is called donor and the individual who receives the graft is called host or recipient.

If the graft is placed into its normal anatomic location, the procedure is called orthotopic transplantation. If the graft is placed in a different site it is called heterotopic transplantation.

Transplantation is the only form of treatment for most end-stage organ failure. In clinical practice, transplantation is used to overcome a functional and anatomic deficit in the recipient. Transplantation of kidneys, hearts, livers, lungs, pancreas and bone marrow are widely done today.

Methods of Transplantation

- Auto grafting: The transfer of self tissue from one body site to another in the same individual
- Allografting: The transfer of organs or tissues from human to human
- Xenografting: The transfer of tissue from one species to another (Figure 11.11).

TN_GOVT_XII_Micro_Biology_CH11.indd 239 04-03-2019 12:32:03