a. genotype b. phenotype
c. both a & b d. none

9. _____ sequence terminates protein synthesis
   a. UAA b. UAG
   c. UGA d. All the above
10. The principal start codon corresponds to which amino acid?
    a. Valine
    b. arginine
    c. Methionine
    d. Isoleucine
11. Number of nucleoprotein subunit in a prokaryotic ribosome
    a. 2 b. 4
    c. 5 d. 6
12. A deletion occurs that eliminates a single amino acid in a protein. How many base pairs were deleted?
    a. 1 b. 2
    c. 3 d. 4
13. During conjugation plasmids undergo
    a. Theta replication
    b. rolling circle replication
    c. sigma replication
    d. gamma replication
14. If a plasmid is mobilizable but nonconjugative, what function does it lack?
    a. Antibiotic resistance
    b. Fertility
    c. Colicinogenic
    d. Restriction sequences
15. The uptake of naked DNA from the surrounding is known as
    a. Transduction
    b. Conjugation
    c. Transformation
    d. Lysis
16. The F plasmid makes a cell
    a. Donor b. Recipient
    c. Resistant d. None
17. Which of the following is more efficient in transferring chromosomal DNA
    a. F^+cell b. F^-cell
    c. Hfr cell d. R^+cell
18. Which of the following statement is true
    a. Protein is the only gene product
    b. A functional gene product is protein or might also be one of several classes of RNA molecules
    c. Carbohydrate is the only gene product
    d. Lipids are the only gene product
19. DNA is transcribed into
    a. mRNA b. tRNA
    c. sRNA d. hnRNA
20. Which of the following is found as part of all prokaryotic promoters
    a. Pribnow box
    b. Shine dalgarno sequence
    c. AUG sequence
    d. UAG sequence

Answer the following

1. What is the direction of synthesis of RNA?
2. Define coding strand.
3. What parts of a mRNA molecule not translated? Ans. Leader & Introns
4. How many codons could be contained in a four-letter code? Ans 4^4=256

TN_GOVT_XII_Micro_Biology_CH12.indd 288 04-03-2019 10:51:56