Observation;

Filamentous hyphae bearing sporangia were observed.

Results:

Wet mount using lactophenol cotton blue was carried to identify the fungus sample. Hyphae with sporangium bearing sporangiospores were observed. It is likely to be of mucor species.

3. Blood Grouping

Aim: To determine the blood group of the blood sample by the slide agglutination test.

Theory and Principle:

Blood grouping is an essential requirement before blood is transfused from one person to another.It is also useful in settling paternity disputes and medicolegal problems.

Red blood cells contain blood group antigens. Antibodies to the blood group antigens are present in the blood plasma.The antigens are generally determined and are responsible for blood types. When RBCs of a person are mixed with corresponding antiserum, agglutination occurs due to antigen-antibody reactions.

Materials Required

- Blood sample (anticoagulated)
- Sterile cotton
- Sterile lancet
- Clean dry grease free slides or white tile
- Toothpicks
- Marker pen
- Commercially available Anti A sera, Anti B sera and Anti D sera

Procedure

1. Prick the finger under aseptic conditions
2. Place a drop of blood on the slide on each side marked as A, B and D.
3. Add a drop of antiserum A , B and D on A, B and D side respectively.
4. Mix with toothpick using separate toothpicks for each mixture.
5. Wait for 2 mins and observe for clumping reaction if any confirm it by observing under microscope.
6. Interpret the results and report.

XII_Micro_Biology_Practical_Manual.indd 297 15-04-2019 17:13:12